

## Review Article

# Memory and Cognition-Related Neuroplasticity Enhancement by Transcranial Direct Current Stimulation in Rodents: A Systematic Review

Carla Cavaleiro,<sup>1,2,3</sup> João Martins,<sup>1,2,3,4</sup> Joana Gonçalves ,<sup>1,2</sup>  
and Miguel Castelo-Branco ,<sup>1,2,3,4</sup>

<sup>1</sup>Coimbra Institute for Biomedical Imaging and Translational Research (CIBIT), University of Coimbra, Coimbra, Portugal

<sup>2</sup>Institute of Nuclear Sciences Applied to Health (ICNAS), University of Coimbra, Coimbra, Portugal

<sup>3</sup>CNC.IBILI Consortium, University of Coimbra, Coimbra, Portugal

<sup>4</sup>Coimbra Institute for Clinical and Biomedical Research (iCBR), Faculty of Medicine, University of Coimbra, Coimbra, Portugal

Correspondence should be addressed to Joana Gonçalves; [jgoncalves@fmed.uc.pt](mailto:jgoncalves@fmed.uc.pt)  
and Miguel Castelo-Branco; [mcbranco@fmed.uc.pt](mailto:mcbranco@fmed.uc.pt)

Received 6 December 2019; Revised 27 January 2020; Accepted 6 February 2020; Published 26 February 2020

Guest Editor: Luca Marsili

Copyright © 2020 Carla Cavaleiro et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Brain stimulation techniques, including transcranial direct current stimulation (tDCS), were identified as promising therapeutic tools to modulate synaptic plasticity abnormalities and minimize memory and learning deficits in many neuropsychiatric diseases. Here, we revised the effect of tDCS on the modulation of neuroplasticity and cognition in several animal disease models of brain diseases affecting plasticity and cognition. Studies included in this review were searched following the terms (“transcranial direct current stimulation”) AND (mice OR mouse OR animal) and according to the PRISMA statement requirements. Overall, the studies collected suggest that tDCS was able to modulate brain plasticity due to synaptic modifications within the stimulated area. Changes in plasticity-related mechanisms were achieved through induction of long-term potentiation (LTP) and upregulation of neuroplasticity-related proteins, such as *c-fos*, brain-derived neurotrophic factor (BDNF), or N-methyl-D-aspartate receptors (NMDARs). Taken into account all revised studies, tDCS is a safe, easy, and noninvasive brain stimulation technique, therapeutically reliable, and with promising potential to promote cognitive enhancement and neuroplasticity. Since the use of tDCS has increased as a novel therapeutic approach in humans, animal studies are important to better understand its mechanisms as well as to help improve the stimulation protocols and their potential role in different neuropathologies.

## 1. Introduction

Transcranial direct current stimulation (tDCS) is a noninvasive brain stimulation technique that promotes transient polarity-dependent changes in spontaneous neuronal activity. This effect is mediated by the application of constant low-amplitude electrical currents using epicranially positioned electrodes above a specific brain region of interest [1–4]. The therapeutic use of low-amplitude electrical currents has a long historical track. Accordingly, both Greeks and Romans used electric torpedo fishes for migraine treatment, and in the 11<sup>th</sup> century, a similar therapeutic procedure was attempted to handle epilepsy [5]. In the 19<sup>th</sup>

century, the application of galvanic currents was attempted to heal melancholia [6]. Over the years, the scientific community interest in brain stimulation grew, and several noninvasive brain stimulation techniques were developed such as tDCS, deep brain stimulation, or transcranial magnetic stimulation. The epicranial application of direct currents promotes a weak electric field force and produces neuronal membrane potential changes [7, 8]. These alterations occur through sodium and calcium currents [1] modulating spontaneous neuronal activity [2]. The consequent regional neuronal inhibition or excitation depends on the applied current polarity [9, 10]. So, it was overall observed that cathodal currents produce inhibitory effects, and thus hyperpolarization,

whereas anodal currents increase excitability in the form of depolarization [2, 11] (Figure 1).

There is nowadays an ongoing discussion regarding the factors that interfere with tDCS outcomes. The initial brain resting state of each subject [12], his/her baseline performance [13], specific individual variations in brain tissue morphology [14], or even more particular details from the experimental design or stimulation protocol used [15] influence these outcomes. *In vivo* and *in vitro* studies are consensual to demonstrate that tDCS-modulated cortical excitability depends on several stimulation parameters, such as duration and frequency of stimulation [16]; polarity, intensity, and density of the applied current [17, 18]; and electrode size and position in the scalp [18–20]. Despite that, beneficial effects of tDCS in several brain disorders, such as PD [21, 22], depression [23], stroke [24, 25], or autism [26, 27], have been documented, and there is growing evidence proposing tDCS application in multiple other disease conditions affecting cognition and neuroplasticity mechanisms.

Both preclinical and clinical studies have demonstrated therapeutic effects of tDCS. Indeed, in human studies, anodal tDCS applied intermittently in the prefrontal cortex (PFC) during slow-wave sleep period, improved recall of declarative memories (word pairs). The authors correlated these findings with enhancement of slow oscillatory electroencephalogram (EEG) activity (<3 Hz, delta ( $\delta$ ) waves), responsible for neuronal plasticity facilitation [28]. Also, anodal tDCS over dorsolateral prefrontal cortex (DLPFC) improved working memory in PD patients and in major depression patients by boosting cortical excitability [21, 23]. Accordingly, preclinical animal studies reported that cortical anodal tDCS improved spatial memory in both wild type (WT) [29] and the AD rat model [30]. Beneficial effects were also found during the early stage of traumatic brain injury (TBI) [31] and following a pilocarpine-induced *status epilepticus* in normal rats [32]. Moreover, improvements were also reported concerning short-term memory in an animal model of attention deficit hyperactivity disorder (ADHD) [33].

The molecular mechanisms underlying the tDCS-mediated cognitive improvements and neuroplasticity processes have become the focus of recent interest. Accordingly, tDCS modulation over several cognition-related plasticity genes and their signaling pathways has been studied. In this review, we provide a state of the art on the application of different protocols of tDCS in animal models highlighting its effectiveness on neuroplasticity mechanisms and, consequently, their related learning and memory processes. Since the published systematic reviews focused on human application of tDCS, here, we provide a comprehensive revision of the effect of tDCS in *in vivo* rodent models of normal and pathological brain functioning.

## 2. Methods

**2.1. Data Sources and Search.** Studies included in this review were identified by searching PubMed. The search was run until 31 October 2019. The search terms were (“transcranial direct current stimulation”) AND (mice OR mouse OR animal). Articles were firstly assessed based on their abstracts

and titles, aiming to include studies that reported applying tDCS to cognitive impairment in animal models. Simultaneously, the following exclusion criteria were adopted to reject studies: (1) not written in English; (2) performing reviews; (3) in human subjects; (4) *in vitro* models; (5) employing other brain stimulation techniques (e.g., transcranial magnetic stimulation (TMS), deep brain stimulation (DBS), or transcranial alternating current stimulation (tACS)); and (6) not explicitly describing the tDCS protocol (stimulation area, number of sessions, frequency, intensity, and pattern).

**2.2. Data Extraction.** A data extraction sheet was developed seeking to retrieve relevant information from each study, namely, study design, sample size, animal model, whether additional therapy was performed, details of the tDCS protocol, outcome measures, and behavioral results.

**2.3. Study Selection.** The database search was elaborated according to the PRISMA statement requirements [34]. 404 records were found, which underwent a preliminary screening (of titles and abstracts), with 314 records being excluded because they did not meet the eligibility criteria. After the full-text analysis of each of the 90 individual articles, 44 rodent studies focusing on tDCS effects over cognition and neuroplasticity in both healthy and neuropathological animal models were selected (Figure 2).

## 3. Results

**3.1. Role of Anodal tDCS in Cognition Processing in Healthy Animals.** In healthy animals, studies demonstrated memory improvement in association with induction of synaptic plasticity mechanisms. In fact, tDCS to prefrontal cortex improved monkey’s performance on an associative learning task by altering low-frequency oscillations and functional connectivity, both locally and between distant brain areas [35]. Regarding rodent models, data are controversial regarding fear condition. Right frontal anodal tDCS administered 24 h before behavioral task did not alter contextual and auditory learning and memory [36]. Additionally, another study described that while the anodal stimulation did not affect fear retrieval, posttraining cathodal stimulation improved fear memory retrieval [37, 38]. However, left prefrontal anodal and cathodal tDCS impaired the acquisition of both contextual and cued fear memory, which could be explained by activity modulation of deep structures such as the amygdala and hippocampus [39].

Concerning learning and memory, de Souza Custódio and colleagues [29] reported better spatial working memory performance following administration of anodal currents to the medial prefrontal cortex (mPFC). In agreement, it was described that administration of hippocampal anodal tDCS improves learning and memory in the Morris water maze and novel object recognition tests [40]. Moreover, memory performance in the passive avoidance learning task was enhanced by anodal stimulation [41]. Also, cortical cathodal stimulation together with visuospatial memory training led to cognitive improvement [42].




FIGURE 1: Illustration of transcranial direct current stimulation in the mice. Anodal stimulation depolarizes the neuronal membrane and enhances excitability. On the other hand, cathodal stimulation hyperpolarizes the neuronal membrane and decreases excitability.




FIGURE 2: Search flow diagram (in accordance with PRISMA statement). Abbreviations: DB: deep brain stimulation; tACS: transcranial alternating current stimulation; TMS: transcranial magnetic stimulation.

The revised *in vivo* animal model studies regarding tDCS effects in memory and cognition of healthy animals are listed below in Table 1.

**3.2. Beneficial Role of tDCS in Brain Diseases.** Overall, reports using animal models of brain diseases described a beneficial role of tDCS in the mitigation of memory symptoms of neurologic conditions such as Alzheimer’s disease (AD) or traumatic brain injury (TBI). More recent studies demonstrated

that tDCS rescued AD-related cognitive symptoms, namely, spatial memory and motor skills [30, 43, 44]. The repetitive stimulation with anodal tDCS in the AD-like dementia rat model reduced the time interval animals needed to reach a food pellet and also decreased the number of errors in the attempt [43]. The same research group showed later that the abovementioned protocol rescued spatial learning and memory in a  $A\beta_{1-40}$ -lesioned AD rat model [30]. Moreover, the impact of tDCS on cognitive performance

TABLE 1: Effect of transcranial direct current stimulation on memory and learning of healthy animals.

Author	Year	Animal model	Specimen; gender	N	Polarity	Position	Stimulation electrode			Stimulation parameters			Reference electrode		Main findings
							Stimulation intensity (mA)	Size (m <sup>2</sup> )	Stimulation duration (min)	Current density (A/m <sup>2</sup> )	Position	Area (cm <sup>2</sup> )	Anesthesia	rtDCS	
Dockery et al. [42]	2011	NDM	Long-Evans rats; males	41	a-tDCS vs. c-tDCS	Frontal cortex (left or right hemisphere)	0.2	0.035	30	57.1	Back	10.5	N	Y (3 days)	↑ Visuospatial working memory (c-tDCS)
de Souza Custódio et al. [29]	2013	NDM	Wistar rats; males	23	a-tDCS	Left mPFC	0.4	0.25	11	N/A	Neck	1	N	Y (5 days)	↑ Spatial working memory (1 h, 4 h, and 10 h poststimulation)
Faraji et al. [96]	2013	NDM	Long-Evans rats; males	24	a-tDCS	Somatosensory cortex (bilateral)	0.065	N/A	10	N/A	Back of skull	N/A	N	Y	↑ Cortical neural density ↑ Motor learning (a-tDCS applied bilaterally or into paw preferred to reaching contralateral hemisphere)
Podda et al. [40]	2016	NDM	C57BL/6 mice; males	16	a-tDCS vs. c-tDCS	Left parietal cortex (dorsal to hippocampal formation)	0.35	0.06	20	N/A	Ventral thorax	5.2	N	N (single session)	↑ Spatial learning and memory (a-tDCS; benefits observable one week after) ↑ BDNF expressions in the <i>hippocampus</i> CREB/CBP pathway activation
Manteghi et al. [36]	2017	NDM	NMRI mice; males	64	a-tDCS	Right frontal cortex	0.2	0.04	20	N/A	Chest	9.5	N/A	N (single session)	↓ Freezing time % and ↑ latency to the freezing (tDCS following 0.1 mg/kg ACPA injection)
Nasehi et al. [37]	2017	NDM	NMRI mice; males	128	a-tDCS vs. c-tDCS	Right frontal cortex	0.2	0.04	20	N/A	Ventral thorax	9.5	N	Y (2 sessions)	↑ Fear memory retrieval/freezing time (a-tDCS and propranolol injection before conditioning)
Nasehi et al. [38]	2017	NDM	NMRI mice; males	128	a-tDCS vs. c-tDCS	Left frontal cortex	0.2	0.04	20	N/A	Ventral thorax	9.5	N	Y (2 sessions)	↑ Contextual fear memory acquisition (a-tDCS before pre- or posttraining)
Abbasi et al. [39]	2017	NDM	NMRI mice; males	41	a-tDCS vs. c-tDCS	Left PFC	0.2	0.04	20 or 30	N/A	Chest	9.5	N	N (single session)	↓ Contextual and cued fear memory

TABLE 1: Continued.

Author	Year	Animal model	Specimen; gender	N	Stimulation electrode			Stimulation parameters			Reference electrode		Main findings		
					Polarity	Position	Stimulation intensity (mA)	Size (m <sup>2</sup> )	Stimulation duration (min)	Current density (A/m <sup>2</sup> )	Position	Area (cm <sup>2</sup> )		Anesthesia	rtDCS
Martins et al. [97]	2019	NDM	Male Wistar rats; males	50	a-tDCS	Left mPFC	0.4	N/A	13	N/A	N/A	N/A	N/A	Y (5 days)	↑ Spatial working memory ↑ GAP-43 (extinct by AMPAR antagonist PRP)
Yu et al. [41]	2019	NDM	Sprague Dawley rats; males	224	a-tDCS	SC, dorsal to the hippocampus	0.25	0.25	30	N/A	N/A	Anterior chest	N/A (EEG electrode)	Y	N (single session) ↑ LTP in CA1 <i>hippocampus</i> (blocked by TrkB antagonist) ↑ BDNF in CA1 <i>hippocampus</i>

Abbreviations: rtDCS: repetitive transcranial direct current stimulation; a-tDCS: anodal transcranial direct current stimulation; c-tDCS: cathodal transcranial direct current stimulation; SC: stereotaxic coordinates; C57BL/6: mouse strain; NMR1: Naval Medical Research Institute outbred mice; NBM: no disease model; SHR: spontaneous hypertensive rats; WKY: Wistar Kyoto rats; PFC: prefrontal cortex; mPFC: medial prefrontal cortex; ITC: inferotemporal cortex; CA1: cornu ammonis 1 region in the hippocampus; PRP: perampanel; ACPA: anticitullinated protein antibody (selective cannabinoid CBI receptor agonist); AMPAR:  $\alpha$ -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid receptor; Trk: tropomyosin receptor kinase receptor; CREB/CBP: cAMP response element binding protein; BDNF: brain-derived neurotrophic factor; GAP-43: growth-associated protein 43; LTP: long-term potentiation; EEG: electroencephalography; A/m<sup>2</sup>: ampere per square meter; mA: milliamperes; cm<sup>2</sup>: square centimeter; mm: millimeter; h: hour; min: minute; vs.: versus; Y: yes; N: no; N/A: not available.

of streptozotocin-induced diabetic rats has been evaluated. Both anodal and cathodal stimulations in the prefrontal cortex restored memory impairment [45, 46] together with restoration of LTP [45]. Other authors evaluated the potential therapeutic effects of tDCS in memory impairment in an animal model of ADHD. It was found that this neuromodulation technique was able to improve short- and long-term memory deficits in the spontaneous hypertensive rats (SHR) but not in their control, Wistar Kyoto rats [33, 47]. In addition, no changes were detected in working memory of these control rats following administration of tDCS [47].

Anodal tDCS also ameliorated behavioral and spatial memory function in the early phase after TBI when it was delivered two weeks postinjury. However, earlier stimulation only improved spatial memory [31]. In a later phase of TBI, it was possible to observe motor recovery as well as spatial memory improvement following repeated anodal tDCS [48]. A growing number of studies has been reporting promising effects of neurostimulation in models of addictive disorders, by reducing craving and maladaptive pervasive learning [49]. In fact, repeated anodal stimulation in mouse frontal cortex decreased nicotine-induced conditioned place preference and further improved working memory [50]. Same polarity currents also could prevent cocaine-induced locomotor hyperactivity and place preference conditioning [51]. In addition, it has been reported that cathodal stimulation has an anticonvulsive effect [16, 32, 52–54]. Indeed, the administration of hippocampal tDCS rescued cognitive impairment by reducing hippocampal neural death and supragranular and CA3 mossy fiber sprouting in a *lithium-pilocarpine-induced status epilepticus* rat [32]. Other neuroplastic effects were evidenced in the reversion of motor symptoms in PD by tDCS administration. The application of anodal currents enhanced graft survival and dopaminergic re-innervation of the surrounding striatal tissue and pronounced behavioral recovery [55].

Despite the fact that many studies reported recovery from memory deficits following tDCS stimulation, there are some opposing reports in animal models of disease affecting cognition. In a recent study from Gondard and collaborators using a triple transgenic (3xTg) mouse model of AD, it was evidenced that a neurostimulation was not able to ameliorate memory symptoms [56]. To reconcile this discrepancy, previous authors have suggested the importance of choosing an optimal current intensity in order to modulate cortical excitability since LTP alterations were dependent on current intensity [57].

The reports regarding tDCS effects in cognition and memory in animal models of brain disease are listed in Tables 2 and 3.

**3.3. Effect of tDCS on Cellular and Molecular Neuroplasticity Mechanisms.** Neuronal network reorganization underlies neuroplasticity processes like developmental synaptogenesis, or neurogenesis and synaptic turnover later on, which ultimately contributes to optimal brain development and aging, as well as functional recovery upon trauma [58]. Interestingly, several reports using genetic engineered animals, phar-

macologically induced animal models of disease, or *in vitro* techniques enlightened the potential of direct current stimulation (DCS) to interact with a myriad of neuroplasticity-related processes such as neuroinflammation [59, 60], neural stem cell migration [59], neurite growth [61], or neurogenesis [62]. Moreover, both human and *in vivo* animal studies evidenced a tDCS-induced effect on memory and learning [28, 35, 63]. However, the underlying cellular and molecular mechanisms remain to be elucidated.

**3.3.1. Modulation of the Excitatory/Inhibitory Network.** To date, animal experimental evidence highlighted tDCS influences on synaptic plasticity, through alterations in the functional connectivity of cognition-related areas [35] and by modulation of excitatory/inhibitory network *tonus* [64], which may involve both the GABAergic and glutamatergic systems. Accordingly, a study conducted with older adults remarked an anodal stimulation effect in gamma-aminobutyric acid (GABA) levels [65]. Similarly, in human healthy volunteers, an anodal tDCS effect in motor learning was correlated with a decrease in GABA levels, an outcome known to be a determinant factor in the promotion of long-term potentiation- (LTP-) dependent plasticity and therefore learning [66, 67].

Several preclinical studies probed LTP enhancement following direct current stimulation. Anodal DCS enhanced LTP in both mouse cortex [68] and rat hippocampal slices [69, 70]. Further, this neurostimulation method increased local field potential (LFPs) amplitudes in primary somatosensory cortex of rabbits [63]. Also, other works demonstrated that neurostimulation-enhanced hippocampal LTP was associated with better spatial memory performance along with an increase in brain-derived neurotrophic factor (BDNF) expression levels [40]. An opposite effect on LTP and LFPs was obtained with administration of cathodal currents. In agreement, a report from Sun et al. [71] evidenced that cathodal currents applied in mouse neocortical slices induced field excitatory postsynaptic potential depression. This type of LTD was smothered by application of an mGluR5 negative allosteric modulator [72]. These findings support a possible modulatory effect of tDCS on mGluR5-mTOR signaling [72]; these molecular pathways are recognized to disturb cognition-related synaptic plasticity.

Further evidence supporting tDCS effect on LTP-like mechanisms was recently brought to light by Stafford et al. [73]. These authors observed that a single anodal tDCS increased both the phosphorylation at the S831 of GluA1 subunit and the translocation of  $\alpha$ -amino-3-hydroxy-5-methyl-4-isoxazole propionic acid receptors (AMPA) from cytosolic to synaptic fractions in the hippocampus. These data could be favoring learning enhancement, as this translocation has been associated with hippocampal LTP induction [72]. Accordingly, others reported a spatial working memory enhancement after anodal stimulation over left medial PFC that was lost with the administration of the AMPAR antagonist perampanel (PRP). In contrast to cathodal currents, anodal currents enhanced intracellular calcium ( $\text{Ca}^{2+}$ ) intake in cell cultures including astrocytes [74–76], a process associated with AMPAR phosphorylation



TABLE 2: Continued.

Author	Year	Animal model	Specimen; gender	N	Polarity	Stimulation electrode			Stimulation parameters			Reference electrode		Main findings		
						Position	Stimulation intensity (mA)	Size (cm <sup>2</sup> )	Stimulation duration (min)	Current density (A/m <sup>2</sup> )	Position	Area (cm <sup>2</sup> )				
Leffa et al. [33]	2016	SHR	SHR rats and WKT rats; males	48	a-tDCS	Frontal cortex	0.5	1.5	20	33.4	Between ears	1.5	N	Y (8 consecutive days)	rtDCS	↑ DA levels in STR in both rat strains and in the hippocampus following tDCS treatment in WKY ↑ BDNF levels in WKY rats Short-term memory improvement
Wu et al. [45]	2017	STZ-induced diabetic rats	Sprague Dawley rats; males	130	a-tDCS	dPFC	0.2	0.0314	30	N/A	Anterior chest	0.25	Y	Y		↑ Spatial working memory and mPFC LTP restoring
Pedron et al. [50]	2017	Cocaine injections	Swiss mice; females	165	a-tDCS	Left frontal cortex	0.2	0.035	2 × 20	N/A	Ventral thorax	9.5	N	Y (5 days stimulation, twice a day; 5 h intersimulation interval)		↓ Cocaine-induced locomotor activity No cocaine-induced place preference (5 mg/kg and 25 mg/kg) ↑ Zif268 basal expression under the electrode area, in the STR and cortex
Leffa et al. [55]	2018	ADHD	SHR and WKY rats; males	30	a-tDCS (bicephalic)	Frontal cortex (supraorbital area)	0.5	1.5 (ECG electrode)	20	33.3	Neck	1.5 (ECG electrode)	N/A	Y (8 days)		↓ Inflammatory cytokines and reversion of long-term memory deficits in SHR rats
Bragina et al. [47]	2018	CCI	Mice; N/A	40	a-tDCS	Parietal somatosensory cortex	0.1	N/A	15	N/A	Ventral thorax	N/A	N/A	Y (daily for 4 days, over 4 weeks and 3 days interval)		Motor coordination recovery Spatial memory and learning performance improvement ↑ CBF bilaterally (regional arteriolar dilatation and hypoxia reduction)
Roostaei et al. [44]	2019	STZ-induced diabetic rats	Wistar rats; males	64	a-tDCS vs. c-tDCS	Left frontal cortex	0.2	3.5	20	N/A	Ventral thorax	9.5	N	Y (twice a day over 2 days)		Restoration of STZ-induced amnesia (both polarities)



TABLE 2: Continued.

Author	Year	Animal model	Specimen; gender	N	Polarity	Position	Stimulation electrode			Stimulation parameters			Reference electrode	Anesthesia	rtDCS	Main findings
							Stimulation intensity (mA)	Size (cm <sup>2</sup> )	Stimulation duration (min)	Current density (A/m <sup>2</sup> )	Position	Area (cm <sup>2</sup> )				
Gondard et al. [55]	2019	Animal model of AD-triple transgenic (3xTg) mice	Triple transgenic (3xTg) mice; males	27	c-tDCS and a-tDCS	Secondary motor cortex (M2) (c-tDCS) and dorsal temporal <i>hippocampus</i> (a-tDCS)	0.05	0.0325	20	N/A	N/A	0.0325	N/A	N/A	Y (5 days/week for 3 weeks)	No treatment effect on memory outcome or AD neuropathological biomarkers

Abbreviations: rtDCS: repetitive transcranial direct current stimulation; a-tDCS: anodal transcranial direct current stimulation; c-tDCS: cathodal transcranial direct current stimulation; TBI: traumatic brain injury; ADHD: attention deficit hyperactivity disorder; SHR: spontaneous hypertensive rats; WKY: Wistar Kyoto rats; CBF: cerebral blood flow; PFC: prefrontal cortex; mPFC: medial prefrontal cortex; dPFC: dorsolateral prefrontal cortex; DG: dentate gyrus; STR: striatum; M2: secondary motor cortex; ITC: inferotemporal cortex; CA3: cornu ammonis 3 region in the *hippocampus*; STZ: streptozotocin; AD: Alzheimer's disease; Aβ<sub>1-40</sub>: amyloid beta peptide 1–40; ACh: acetylcholine; BDNF: brain-derived neurotrophic factor; DA: dopamine; GFAP: glial fibrillary acidic protein; Zlf268: zinc finger transcription factor 268; LTP: long-term potentiation; IP: intraperitoneal; SC: subcutaneous; SE: *status epilepticus*; ECG: electrocardiography; mg: milligram; kg: kilogram; A/m<sup>2</sup>: ampere per square meter; mA: milliampere; cm<sup>2</sup>: square centimeter; mm: millimeter; h: hour; min: minute; vs.: versus; Y: yes; N: no; N/A: not available.

TABLE 3: Role of transcranial direct current stimulation on neuroplasticity with a focus on animal models of neurotrauma.

Author	Year	Animal model	Specimen; gender	N	Stimulation parameters					Main findings					
					Polarity	Position	Stimulation intensity (mA)	Stimulation electrode Size (cm <sup>2</sup> )	Stimulation duration (min)		Current density (A/m <sup>2</sup> )	Reference electrode Position	Area (cm <sup>2</sup> )	Anesthesia	rtDCS
Nekhendzy et al. [97]	2004	Inflammatory nociception model	Sprague Dawley rats; males	31	c-tDCS	Frontal cortex	2.25	N/A	45	N/A	Bimastoid	N/A	Y	Y (8 days)	↓ Nociceptive response (effects lasted up to 50 min poststimulation)
Taib et al. [98]	2009	Left hemispherectomy	Wistar rats; males	9	a-tDCS	Right or left motor cortex	0.4	0.071	20	56.3	Supraorbital region	0.0064	N/A	Y	↑ T corticomuscular response amplitudes
Kim et al. [99]	2010	MCAO	Sprague Dawley rats; both genders	61	c-tDCS vs. a-tDCS	Left primary motor cortex (M1)	0.1	0.785	30	N/A	Trunk	9	Y	Y (2 weeks)	Neuroprotection over white matter and ischemic size ↓ (a-tDCS) ↑ Motor function
Jiang et al. [95]	2012	MCAO	Sprague Dawley rats; both genders	90	a-tDCS and c-tDCS	Motor cortex	0.1	0.785	30	1.27	Trunk	9	Y	Y (daily, for 3, 7, or 14 days postlesion)	↑ Motor function (7 and 14 days poststroke) ↑ Dendritic spine density ↓ PXI expression (7th and 14h day poststroke)
Yoon et al. [56]	2012	MCAO	Sprague Dawley rats; males	30	a-tDCS	Left primary motor cortex (M1)	0.2	N/A	20	28.2	Anterior chest	48.0	Y	Y (5 days stimulation 1 week vs. 1 day postischemia)	↑ Motor function and Barnes maze performance (a-tDCS applied 1 week after ischemic injury) ↑ MAP-2 and GAP-43 expression around the perilesional area (a-tDCS applied 1 week postischemic injury)
Laste et al. [100]	2012	CFA injection/chronic inflammation induction	Wistar rats; males	18	a-tDCS	Parietal cortex	0.5	1.5 (ECG electrode)	20	33.4	Supraorbital area	1.5 (ECG electrode)	N	Y (8 days)	Significant differences in nociceptive response (immediately after and 24 h poststimulation)
Adachi et al. [101]	2012	CRS	Wistar rats; males	48	a-tDCS	Parietal cortex	0.5	1.5 (ECG electrode)	20	3.3	Supraorbital area	1.5 (ECG electrode)	N	Y (8 days)	↓ Nociceptive response in chronic stress condition ↓ TNF expression in the hippocampus

TABLE 3: Continued.

Author	Year	Animal model	Specimen; gender	N	Polarity	Position	Stimulation electrode			Stimulation parameters			Reference electrode Position	Reference electrode Area (cm <sup>2</sup> )	Anesthesia	rtDCS	Main findings
							Stimulation intensity (mA)	Size (cm <sup>2</sup> )	Stimulation duration (min)	Current density (A/m <sup>2</sup> )							
Peruzzotti-Jametti et al. [60]	2013	MCAO	C57BL/6 mice; males	137	c-tDCS vs. a-tDCS	Left parietal cortex	0.25	0.0144	40	55	Ventral thorax	5.2	N/A	N (single session)	↑ Infarct volume and ↑ BBB leakage (a-tDCS ipsilesional hemisphere) Cortical Glu activity ↓, functional ↑ and ischemic damage ↓ (c-tDCS ipsilesional hemisphere)		
Notturmo et al. [102]	2014	MCAO	Sprague Dawley rats; both genders	53	c-tDCS	Left motor cortex	0.2	0.07	120	28.6	Ventral thorax	10.5	Y	N (single session)	Ischemia volume ↓		
Lu et al. [103]	2015	MPTP injection	C57bl mice; males	36	a-tDCS	Left frontal cortex	0.2	0.035	10	57	Between shoulders	9	N/A	Y (daily for 3 weeks)	↑ Motor coordination (until 21 days poststimulation) ↑ TH and DA expression ↓ Oxidative stress		
Spezia Adachi et al. [104]	2015	Restraint stress model	Wistar rats; males	78	a-tDCS	Parietal cortex (midline)	0.5	1.5 (ECG electrode)	20	N/A	Supraorbital area	1.5 (ECG electrode)	N	Y (daily for 8 days)	↓ Stress-induced nociceptive response ↑ Pain threshold ↓ BDNF levels (spinal cord and brainstem) in unstressed animals		
Yoon et al. [31]	2016	Lateral fluid percussion method	Sprague Dawley rats; males	36	a-tDCS	Hippocampus	0.2	0.0225	20	28.2	Chest	48 (corset)	Y	Y (daily for 5 days)	↑ Perilesional area BDNF expression (tDCS 2 weeks post-TBI) ↑ Choline/creatinine ratios (tDCS 1 week post-TBI) Motor performance recovery (2 weeks of tDCS)		
Leffa et al. [33]	2016	SHR	SHR rats and WKT rats; males	48	a-tDCS	Frontal cortex	0.5	1.5	20	33.4	Between ears	1.5	N	Y (8 consecutive days)	↑ DA levels in STR in both rat strains and in the hippocampus following tDCS treatment in WKY rats ↑ BDNF levels in WKY rats		

TABLE 3: Continued.

Author	Year	Animal model	Specimen; gender	N	Polarity	Position	Stimulation electrode			Stimulation parameters			Reference electrode	Anesthesia	rtDCS	Main findings
							Stimulation intensity (mA)	Size (cm <sup>2</sup> )	Stimulation duration (min)	Current density (A/m <sup>2</sup> )	Position	Area (cm <sup>2</sup> )				
Liu et al. [105]	2016	PTI	Sprague Dawley rats; N/A	58	c-tDCS	Right S1FL	2	N/A	20	20.37	Venter	N/A	N/A	N (single session)	Ischemia expansion inhibition (c-tDCS immediately posts ischemia induction) ↓ NeuN expression (c-tDCS+PSS group)	
Braun et al. [106]	2016	MCAO	Wistar rats; males	41	c-tDCS vs. a-tDCS	Left primary motor cortex (M1)	0.5	0.035	15	N/A	Ventral thorax	N/A	Y	Y (10-day stimulation; 5 days with 2-day interval)	Gait recovery at day 16 poststroke (both polarities) Faster recovery of limb strength (fully recovered strength at day 10 and gait at day 14 with c-tDCS) ↑ Microglia and neuroblasts in lesion ipsilateral cortex	
Cioato et al. [107]	2016	Sciatic nerve chronic constriction	Wistar rats; males	84	a-tDCS and c-tDCS	Parietal cortex (bicephalic)	0.5	1.5 (ECG electrode)	20	N/A	Supraorbital area	1.5 (ECG electrode)	N	Y (8 days)	Nociceptive relieve (for up to 7 days poststimulation) Reversion of T IL-1 $\beta$ levels (48 h and 7 days poststimulation)	
Filho et al. [108]	2016	Partial sciatic nerve compression	Wistar rats; males	144	a-tDCS	Parietal cortex	0.5	1.5 (ECG electrode)	20	N/A	Supraorbital area	1.5 (ECG electrode)	N	Y (8 days)	↓ BDNF expression (48 h poststimulation) Reversion of behavioral alterations (analgesic and anxiolytic) associated with neuropathic pain	
Moreira et al. [109]	2016	Pain and menopause (ovariectomised animals)	Wistar rats; females	45	C-tDCS	Parietal cortex	0.5	1.5 (ECG electrode)	20	N/A	Supraorbital area	1.5 (ECG electrode)	N	Y (8 days)	↓ Hypothalamic BDNF levels and T serum BDNF in ovariectomised animals	
Dimov et al. [87]	2016	N/A	Wistar rats; males	25	c-tDCS	Left primary motor cortex (M1)	0.25	0.0227	15	N/A	Ventral thorax	N/A	N	N (single session)	Bilateral ↓ Egr-1 expression in the PAG Spinal ENK immunoreactivity ↓ in the DHSC	

TABLE 3: Continued.

Author	Year	Animal model	Specimen; gender	N	Stimulation parameters				Main findings						
					Polarity	Position	Stimulation intensity (mA)	Stimulation electrode Size (cm <sup>2</sup> )		Stimulation duration (min)	Current density (A/m <sup>2</sup> )	Reference electrode Position	Area (cm <sup>2</sup> )	Anesthesia	rtDCS
Liu et al. [110]	2017	PTI	Sprague Dawley rats; males	58	c-tDCS	S1FL	2	N/A	20	N/A	3 mm lateral to lambda	N/A	N/A	N (single session)	Prevention of ischemia injury expansion during hyperacute phase of ischemia (c-tDCS+PSS)
Kim & Han [111]	2017	Modified Tang's method [128]	Sprague Dawley rats; N/A	31	a-tDCS	Left motor cortex	0.2	1	30	0.26	Ventral thorax	9	Y	N	Early recovery of consciousness and MEP and SEP prolonged latency (tDCS applied right after TBI) ↓ Astroglial GFAP immunoreactivity
Winkler et al. [112]	2017	Striatal 6-OHDA injection	Sprague Dawley rats; females	24	a-tDCS and c-tDCS	Left motor cortex	N/A	0.16	20	8	Chest	3	N	Y (daily for 14 days)	Graft survival, striatal dopaminergic reinnervation and motor recovery (a-tDCS)
de Souza et al. [113]	2017	PSNL	Swiss mice; males	N/A	a-tDCS and c-tDCS	Parietal cortex (bicephalic)	0.5	N/A (EEG electrode)	5; 10; 15; 20	N/A	Supraorbital area	N/A (EEG electrode)	N	N (single session)	Antiallodynamic effect (seen 4 h poststimulation of 15 min and 20 min)
Leffa et al. [46]	2018	ADHD	SHR and WKY rats; males	30	a-tDCS (bicephalic)	Frontal cortex (supraorbital area)	0.5	1.5 (EEG electrode)	20	33.3	Neck	1.5 (EEG electrode)	N/A	Y (8 days)	↓ Inflammatory cytokines and reversal of long-term memory deficits in SHR rats
Paciello et al. [114]	2018	NIHL	Wistar rats; males	124	a-tDCS	Temporal lobe (auditory cortex)	0.35	0.0625	20	56	Ventral thorax	12	N	Y (2 days)	↑ Dendritic spines density (layer 2/3 pyramidal neurons of the auditory cortex) ↑ BDNF and synaptophysin expression in auditory cortex (24 h poststimulation)
Fregni et al. [115]	2018	N/A	Wistar rats; males	32	N/A	N/A (bicephalic)	N/A	N/A	20	N/A	N/A	N/A	N/A	Y (8 days)	tDCS prior to stress exposure prevented thermal hyperalgesia

TABLE 3: Continued.

Author	Year	Animal model	Specimen; gender	N	Polarity	Position	Stimulation electrode		Stimulation parameters			Reference electrode Position	Reference electrode Area (cm <sup>2</sup> )	Anesthesia	rtDCS	Main findings
							Stimulation intensity (mA)	Size (cm <sup>2</sup> )	Stimulation duration (min)	Current density (A/m <sup>2</sup> )						
Lee et al. [116]	2019	MPTP injection	C57bl mice; male	60	a-tDCS	Primary motor cortex (M1)	N/A	N/A	30	N/A	Between shoulders	N/A	N/A	Y (daily for 5 days)	<p>↑ Motor coordination</p> <p>Rescue of MPTP-induced mitochondrial dysfunction (C, ATP and GDH and \$ Drp1 levels)</p>	
Callai et al. [117]	2019	CCI-ION	Wistar rats, males	151	a-tDCS	Parietal cortex (bicephalic)	0.5	1.5 (ECG electrode)	20	N/A	Supraorbital area	1.5 (ECG electrode)	N	Y (8 days)	<p>↓ Mechanical hyperalgesia</p> <p>↓ TNF-<math>\alpha</math> expression (7 days poststimulation)</p> <p>↓ IL-10 (7 days poststimulation)</p> <p>↓ LDH serum levels</p>	
Scarabelot et al. [118]	2019	CFA injection/chronic inflammation induction	Sprague Dawley rats; males	104	a-tDCS	Parietal cortex (bicephalic)	0.5	1.5 (ECG electrode)	20	N/A	Supraorbital area	1.5 (ECG electrode)	N/A	Y (8 days)	<p>↓ Thermal and mechanical hyperalgesia</p> <p>↑ IL-6 (in brainstem 24 h poststimulation)</p> <p>↓ IL-10 (7 days poststimulation)</p> <p>Normalization of BDNF levels (24 h poststimulation)</p>	

Abbreviations: rtDCS: repetitive transcranial direct current stimulation; a-tDCS: anodal transcranial direct current stimulation; c-tDCS: cathodal transcranial direct current stimulation; C57BL/6: mouse strain; SHR: spontaneous hypertensive rats; WKY: Wistar Kyoto rats; ADHD: attention deficit hyperactivity disorder; 6-OHDA: 6-hydroxydopamine; MPTP: 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine; TBI: traumatic brain injury; PTI: photothrombic ischemia; MCAO: middle cerebral artery occlusion; PSS: peripheral sensory stimulation; NIH1: noise-induced hearing loss; CC-ION: chronic constriction of the infraorbital nerve (pain model); PSNL: partial sciatic nerve ligation (pain model); CRS: chronic restraint stress (pain model); CFA: complete Freund's adjuvant (pain model); BBB: blood-brain barrier; SJFL: forelimb region of the primary somatosensory cortex; M1: primary motor area; PAG: periaqueductal grey; DHSC: dorsal horn of the spinal cord; MEP: motor-evoked potentials; SEP: somatosensory evoked potentials; GFAP: glial fibrillary acidic protein; BDNF: brain-derived neurotrophic factor; Glu: glutamate NeuN: neuronal marker; PX1: pannexin 1; TH: thyroxine hydroxylase; Egr-1: early growth response protein 1; TNF: tumor necrosis factor; IL-1 $\beta$ : interleukin 1 beta; IL-6: interleukin 6; IL-10: interleukin 10; LDH: lactate dehydrogenase enzyme; ATP: adenosine triphosphate; GDH: glutamate dehydrogenase; Drp1: dynamin-related protein; MAP-2: microtubule associated protein 2; GAP-43: growth associated protein 43; ENK: early embryo specific NK; DA: dopamine; ECG: electrocardiography; EEG: electroencephalogram; A/m<sup>2</sup>: ampere per square meter; mA: milliampere; vs.: versus; min: minute; cm<sup>2</sup>: square centimeter; mm: millimeter; Y: yes; N: no; N/A: not available.

and trafficking to postsynaptic density [77] and ultimately, allowing LTP facilitation, a cellular correlate of learning and memory.

**3.3.2. Activation of Neuroplasticity-Associated Gene Expression.** Neurostimulation could have long-lasting effects in memory as data from different studies evidenced [40]. Authors have been argued that tDCS cognition modulation is associated with neuroplasticity-associated gene expression alterations [78]. One of the neuroplasticity-associated genes, known to be essential for hippocampal LTP, is BDNF [79]. Several studies elucidated the role of BDNF in memory modulation by tDCS. In fact, it was reported that anodal currents could increase BDNF expression [68], and its activation *via* tropomyosin receptor kinase (Trk) receptors [80], triggering NMDAR opening, and inducing a later phase LTP (L-LTP) facilitation [81]. Accordingly, Yu et al. [41] found that the administration the Trk inhibitor ANA-12 prevented the anodal tDCS-induced hippocampal CA1 LTP increase. Other studies, using the same polarity currents, revealed a link between the upregulation of BDNF and cAMP response element binding protein/CREB-binding protein (CREB/CBP) [40] involved in LTP and memory formation [82]. Also, the application of cortical anodal currents in frontal cortex was able to upregulate BDNF together with striatal dopamine [33]. The upregulation of BDNF following neurostimulation was associated with the augmentation of expression levels of immediate early genes (IEGs), such as *c-fos* and *zif268* [69]. Moreover, Kim et al. [78] confirmed that repetitive anodal tDCS in right sensorimotor cortex of healthy rats promoted a significant increase of mRNA levels of plasticity-associated genes, namely, BDNF, cAMP response element binding protein (CREB), synapsin I, Ca<sup>2+</sup>/calmodulin-dependent protein kinase II (CaMKII), activity-regulated cytoskeleton-associated protein (Arc), and *c-fos*. It was also demonstrated that sensory evoked cortical responses were boosted after tDCS *via* alpha-1 adrenergic receptor-mediated astrocytic Ca<sup>2+</sup>/IP3 signaling, thus involving also glia and the adrenergic system [75]. Anodal tDCS actions in glia were further confirmed by Mishima et al. [76]. Using a mouse model lacking Ca<sup>2+</sup> uptake in astrocytes, the inositol trisphosphate receptor type 2 (IP3R2) knockout (KO) mouse and also an adrenergic receptor antagonist, they confirmed decreased microglia motility along with soma enlargement in tDCS stimulated animals [76].

In poststroke recovery, it was found that anodal currents significantly increased the GAP-43 and the microtubule-associated protein 2 (MAP-2) expression around the infarct area [56]. These neuronal growth-promoting proteins are overexpressed during dendritic remodeling and axonal regrowth throughout the acute phase of stroke [83, 84]. Anodal stimulation also modulated pannexin-1 (PX1) hemichannel levels [85, 86] and, following an ischemic insult, neurostimulation decreased rat PX1 mRNA and, consequently, augmented dendritic spine density in the surrounding areas of cerebral infarction; these cellular outcomes were associated with the improvement of motor function [85]. Some authors proposed that tDCS-induced improvement of stroke/TBI symptoms might be due to increase of BDNF expres-

sion and associated with choline/creatine ratios in the perilesional cortex [31].

Overall, tDCS methodology was able to modulate molecular pathways involved in the regulation of cognition-related synaptic plasticity mechanisms (Figure 3). The revised *in vivo* animal studies regarding tDCS-induced effects in the cellular and molecular mechanisms of memory and learning are listed in Table 4.

## 4. Discussion

This systematic review collected several studies that confirm the potential effects of tDCS on neuronal activity and synaptic plasticity. Here, we documented a variable combination of stimulation protocols, stimulation areas, and healthy and disease animal models. Most of the existent literature is focused on human application of tDCS. The comprehensive revision of the effect of tDCS on rodent models of normal and pathological brain functioning does therefore provide a novel contribution to the field. Overall, the revised studies indicated that tDCS was able to modulate synaptic plasticity and, consequently, learning and memory processes [87, 88].

Memory formation and consolidation are recognized to rely on activity-dependent modifications, such as LTD and LTP [89], both dependent on the activation of calcium-dependent kinases (e.g., CaMKs), which in turn control the trafficking of NMDARs and AMPARs [90]. Despite the wide set of stimulation protocols, tDCS-induced modulation of NMDAR signaling and synaptic protein upregulation resulting in LTP and cognitive enhancement have been consistently reported in animal studies. Anodal tDCS increased AMPAR synapse translocation [73, 89] and induced spatial memory improvement by involving both CREB and BDNF expression alterations [53]. Also, an increase in hippocampal and cortical mRNA levels of *c-fos*, synapsin, CaMKII, and Arc was observed poststimulation [78].

Similar results highlighting tDCS effects in neuroplasticity were obtained with *in vitro* studies. Accordingly, Ranieri and coworkers [69] probed that anodal currents increased NMDAR-dependent LTP in hippocampal CA3-CA1 synapses [69], in part, due to production of BDNF [68]. In addition, it was demonstrated that tDCS-induced hippocampal BDNF release is dependent on histone acetylation of BDNF gene promoters [40]. Overall, the abovementioned works provide positive evidence for the effect of tDCS on cognitive function enhancement.

Although tDCS impaired the acquisition of both contextual and cued fear memory [39], there are no studies on possible cascades/proteins involved in tDCS-induced neuroplasticity alterations following fear memory changes. Nevertheless, a very recent paper demonstrated chronic repetitive TMS of the ventromedial prefrontal cortex reversed stress-induced behavior impairments with an increase of *c-fos* activity [91].

Cortical anodal currents have been shown to be mostly excitatory and support memory enhancement and neuroplasticity. The literature is also consistent with the notion that the stimulation over the cortical region functionally involved in a certain cognitive task increases performance in that specific task. Marshall et al. demonstrated that anodal




FIGURE 3: Schematic illustration of molecular mechanisms underlying the effect of anodal transcranial direct current stimulation (tDCS) on neuronal physiology. The neurostimulation in the target cortical area depolarizes neuronal membrane and glutamate released in presynaptic neuron and binds in NMDA and AMPA receptors (see book chapter Rozisky et al., 2015). Consequently, there is intracellular  $Ca^{2+}$  upregulation in the postsynaptic neuron, which can activate protein kinases that in turn modulate numerous neuronal signaling pathways (such as the mTOR pathway) leading to transcriptional changes. The tDCS also activates molecular cascades to promote BDNF production. As a long-term mechanism, gene transcription is modulated leading to the formation of new proteins that in turn lead to facilitation of LTP and improvement of cognition. Abbreviations: AMPA:  $\alpha$ -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid; BDNF: brain-derived neurotrophic factor; CBP: CREB-binding protein; CREB: cAMP response element binding protein; GSK3: glycogen synthase kinase 3; LTP: long-term potentiation; mTOR: mammalian target of rapamycin; NMDA: N-methyl-D-aspartate; TrkB: tropomyosin receptor kinase B.

currents over the PFC, a region involved in memory encoding, during slow wave sleep improved declarative memory [28]. However, it was described that cortical cathodal stimulation simultaneously with training task was able to increase visuospatial working memory, in spite of the fact that it was associated with decreased excitability [42]. This suggests that modulatory effects of tDCS were influenced by the polarity-dependent electrical dynamics established between the stimulated area and its related neuronal networks. In agreement, a recent report observed an inhibitory effect in motor learning tasks following anodal currents in the cerebellum; the anodal excitatory effect over the Purkinje cell activity led to an overall inhibition of downstream structures, reducing as a result the vestibulo-ocular reflex gain [90]. Similar paradoxical results have been observed in humans. Recently, Moliadze and collaborators [92] reported that tDCS-induced neural modulation depended on several parameters, namely, the age. In fact, an excitatory effect was seen in young subjects, but not in the older participants.

Nowadays, TMS, another important noninvasive brain stimulation technique, is useful for evaluating excitability in the primary motor cortex (M1) and conductivity along the cortical-spinal tract. This technique has been amply used in rehabilitation of stroke patients [93] and in neuropsychiatric

disorders, namely, depression [94]. tDCS and TMS are undergoing the most active investigation and share a capacity to modulate regional cortical excitability, and both are well-tolerated by children and adults [95]. However, TMS has been already approved for clinical use and tDCS is still undergoing investigation as a plausible therapy for a range of neuropsychiatric disorders [95]. The rationale, in part, for this is because data on the efficacy and safety of tDCS are sparse and employ heterogeneous stimulation protocols. Indeed, there is a paucity of strictly conducted randomized, sham controlled clinical trials, and case considerable follow-up periods, which makes it difficult to use these results to inform clinical practice concerning the putative beneficial role of tDCS. Moreover, tDCS effects seem to be clearly dependent on structure, connectivity, and function of the target brain region. Importantly, these outcomes were intrinsically correlated with GABAergic neurotransmission which raises the issue that one has to take into account that during development GABA can act as an excitatory neurotransmitter [96].

## 5. Conclusions

There is growing evidence that tDCS modulates brain activity and, consequently, enhances synaptic plasticity and cognitive



TABLE 4: Cellular and molecular mechanisms underlying transcranial direct current stimulation effect in the brain.

Author	Year	Animal model	Specimen; gender	N	Stimulation electrode			Stimulation parameters			Reference electrode		Main findings		
					Polarity	Position	Stimulation intensity (mA)	Size (cm <sup>2</sup> )	Stimulation duration (min)	Current density (A/m <sup>2</sup> )	Position	Area (cm <sup>2</sup> )		Anesthesia	rtDCS
Márquez-Ruiz et al. [63]	2012	NDM	New Zealand White albino rabbits; N/A	13	a-tDCS and c-tDCS	Somatosensory cortex (S1)	0.5; 1; 1.5 and 2	0.7857	10	3.7	Ear	35	N	N (single session)	↑ LFP in S1 (a-tDCS) and ↓ LFP S1 (c-tDCS)
Rohan et al. [57]	2015	NDM	Sprague Dawley rats; male	34	a-tDCS	SC dorsal to the <i>hippocampus</i>	0.1 or 0.25	0.25	30	N/A	Between shoulders	8.04	N	N (single session)	# LTP and PPF in the <i>hippocampus</i>
Yoon et al. [31]	2016	Lateral fluid percussion method	Sprague Dawley rats; males	36	a-tDCS	<i>Hippocampus</i>	0.2	0.0225	20	28.2	Chest	48 (corset)	Y	Y (daily for 5 days)	# perilesional area BDNF expression (tDCS 2 weeks post-TBI) # choline/creatinine ratios (tDCS 1 week post-TBI) Motor performance recovery (2 weeks of tDCS)
Leffa et al. [33]	2016	SHR	SHR rats and WKY rats; males	48	a-tDCS	Frontal cortex	0.5	1.5	20	33.4	Between ears	1.5	N	Y (8 consecutive days)	# DA levels in STR in both rat strains and in the hippocampus following tDCS treatment in WKY rats # BDNF levels in WKY rats Short-term memory improvement
Podda et al. [40]	2016	NDM	C57BL/6 mice; males	16	a-tDCS vs. c-tDCS	Left parietal cortex (dorsal to hippocampal formation)	0.35	0.06	20	N/A	Ventral thorax	5.2	N	N (single session)	# spatial learning and memory (a-tDCS); benefits observable one week after # BDNF expression in the <i>hippocampus</i> CREB/CBP pathway activation

TABLE 4: Continued.

Author	Year	Animal model	Specimen; gender	N	Polarity	Position	Stimulation electrode			Stimulation parameters			Reference electrode		Main findings
							Stimulation intensity (mA)	Size (cm <sup>2</sup> )	Stimulation duration (min)	Current density (A/m <sup>2</sup> )	Position	Area (cm <sup>2</sup> )	Anesthesia	rtDCS	
Monai et al. [75]	2016	NDM	G7NG817 mice; N/A	10	a-tDCS	Primary visual cortex (V1)	0.1	0.02	10	N/A	Neck	N/A	N	N (single session)	Up to 50% expansion of visual evoke active area (up to 2 h poststimulation effect) tDCS-induced plasticity depends on the activity of IP, R2, and A1AR
Kim et al. [78]	2017	NDM	Sprague Dawley rats; males	90	a-tDCS	Right sensorimotor cortex	0.25	0.071	20	N/A	Right anterior chest	0.5	N	Y (7 days)	# BDNF, CREB, synapsin, and CaMKII mRNA expression levels (ipsilateral cortex) and c-fos ( <i>hippocampus</i> )
Stafford et al. [73]	2018	NDM	Sprague Dawley rats; males	16	a-tDCS	Caudal to bregma	0.25	0.25	30	N/A	Ventral thorax	N/A	N	N (single session)	↑ AMPAR translocation to the synapse in the hippocampus and ↑ phosphorylation of the S831 site on GluA1
Martins et al. [97]	2019	NDM	Male Wistar rats; males	50	a-tDCS	Left mPFC	0.4	N/A	13	N/A	N/A	N/A	N/A	Y (5 days)	↑ Spatial working memory ↑ GAP-43 (extinct by AMPAR antagonist PRP)
Yu et al. [41]	2019	NDM	Sprague Dawley rats; males	224	a-tDCS	SC dorsal to the <i>hippocampus</i>	0.25	0.25	30	N/A	Anterior chest	N/A (EEG electrode)	Y	N (single session)	↑ Memory (passive avoidance memory retention) ↑ LTP in CA1 <i>hippocampus</i> (blocked by TrkB antagonist) ↑ BDNF in CA1 <i>hippocampus</i>

Abbreviations: rtDCS: repetitive transcranial direct current stimulation; a-tDCS: anodal transcranial direct current stimulation; c-tDCS: cathodal transcranial direct current stimulation; C57BL/6: mouse strain; SC: stereotaxic coordinates; NDM: no disease model; SHR: spontaneous hypertensive rats; WKY: Wistar Kyoto rats; PFC: prefrontal cortex; mPFC: medial prefrontal cortex; dPFC: dorsolateral prefrontal cortex; DG: dentate gyrus; STR: *striatum*; S1: somatosensory cortex; V1: primary visual cortex; ITC: inferotemporal cortex; CA1: cornu ammonis 1 region in the hippocampus; TBI: traumatic brain injury; PRP: perampanel; CREB: cAMP response element-binding protein (transcription factor); CREB/CBP: cAMP response element binding protein; BDNF: brain-derived neurotrophic factor; DA: dopamine; GAP-43: growth associate protein 43; CaMKII: Ca<sup>2+</sup>/calmodulin-dependent protein kinase; mRNA: messenger ribonucleic acid; IP3R2: inositol triphosphate type 2 receptor; A1AR: adenosine A2A receptor; AMPAR: α-amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid receptor; GluA1: AMPA receptor subunit A1; LFP: local field potential; LTP: long-term potentiation; PPF: paired pulse facilitation; EEG: electroencephalography; A/m<sup>2</sup>: ampere per square meter; mA: milliampere; cm<sup>2</sup>: square centimeter; mm: millimeter; h: hour; min: minute; vs.: versus; Y: yes; N: no; N/A: not available.

performance. Overall, reports from laboratory animal research present tDCS as a promising noninvasive brain stimulation technique. The presented evidence is therefore consistent with human studies suggesting that this technique is useful to mitigate neurologic symptoms of several brain disorders, thus improving learning and memory. Further research is needed so that this technique can be fully translated into optimal therapeutic strategies.

### Conflicts of Interest

The authors declare that there is no conflict of interest regarding the publication of this paper.

### Authors' Contributions

Joana Gonçalves and Miguel Castelo-Branco share senior authorship.

### Acknowledgments

This work was supported by grants POCI-01-0145-FEDER-016428 and CENTRO-01-0145-FEDER-000016 financed by Centro 2020 FEDER, COMPETE, FLAD Life Sciences Ed 2 2016, FCT/UID 4950 COMPETE, POCI-01-0145-FEDER-0007440, FCT, and European Grant H2020 STIPED.

### References

- [1] G. Ruffini, F. Wendling, I. Merlet et al., "Transcranial current brain stimulation (tCS): models and technologies," *IEEE Transactions on Neural Systems and Rehabilitation Engineering*, vol. 21, no. 3, pp. 333–345, 2013.
- [2] M. A. Nitsche and W. Paulus, "Excitability changes induced in the human motor cortex by weak transcranial direct current stimulation," *The Journal of Physiology*, vol. 527, no. 3, pp. 633–639, 2000.
- [3] T. Wagner, A. Valero-Cabre, and A. Pascual-Leone, "Noninvasive human brain stimulation," *Annual Review of Biomedical Engineering*, vol. 9, no. 1, pp. 527–565, 2007.
- [4] M. Nitsche, D. Liebetanz, A. Antal, N. Lang, F. Tergau, and W. Paulus, "Modulation of cortical excitability by weak direct current stimulation—technical, safety and functional aspects," *Supplements to Clinical Neurophysiology*, vol. 56, pp. 255–276, 2003.
- [5] P. Kellaway, "The part played by electrical fish in the early history of bioelectricity and electrotherapy," *Bulletin of the History of Medicine*, vol. 20, no. 2, pp. 130–134, 1946.
- [6] A. Parent, "Giovanni Aldini: from animal electricity to human brain stimulation," *The Canadian Journal of Neurological Sciences*, vol. 31, no. 4, pp. 576–584, 2004.
- [7] T. Bullock and C. Terzuolo, "Diverse forms of activity in the somata of spontaneous and integrating ganglion cells," *The Journal of Physiology*, vol. 138, no. 3, pp. 341–364, 1957.
- [8] C. Poreisz, K. Boros, A. Antal, and W. Paulus, "Safety aspects of transcranial direct current stimulation concerning healthy subjects and patients," *Brain Research Bulletin*, vol. 72, no. 4–6, pp. 208–214, 2007.
- [9] L. Bindman, O. Lippold, and J. Redfearn, "The action of brief polarizing currents on the cerebral cortex of the rat (1) during current flow and (2) in the production of long-lasting after-effects," *The Journal of Physiology*, vol. 172, pp. 369–382, 1964.
- [10] C. A. Terzuolo and T. H. Bullock, "Measurement of imposed voltage gradient adequate to modulate neuronal firing," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 42, no. 9, pp. 687–694, 1956.
- [11] M. A. Nitsche and W. Paulus, "Sustained excitability elevations induced by transcranial DC motor cortex stimulation in humans," *Neurology*, vol. 57, no. 10, pp. 1899–1901, 2001.
- [12] A. Antal, D. Terney, C. Poreisz, and W. Paulus, "Towards unravelling task-related modulations of neuroplastic changes induced in the human motor cortex," *The European Journal of Neuroscience*, vol. 26, no. 9, pp. 2687–2691, 2007.
- [13] B. Krause and R. Cohen Kadosh, "Not all brains are created equal: the relevance of individual differences in responsiveness to transcranial electrical stimulation," *Frontiers in Systems Neuroscience*, vol. 8, 2014.
- [14] M. Bikson, A. Rahman, and A. Datta, "Computational models of transcranial direct current stimulation," *Clinical EEG and Neuroscience*, vol. 43, no. 3, pp. 176–183, 2012.
- [15] A. Fertonani and C. Miniussi, "Transcranial electrical stimulation: what we know and do not know about mechanisms," *The Neuroscientist*, vol. 23, no. 2, pp. 109–123, 2017.
- [16] D. Liebetanz, F. Klinker, D. Hering et al., "Anticonvulsant effects of transcranial direct-current stimulation (tDCS) in the rat cortical ramp model of focal epilepsy," *Epilepsia*, vol. 47, no. 7, pp. 1216–1224, 2006.
- [17] D. Liebetanz, R. Koch, S. Mayenfels, F. König, W. Paulus, and M. A. Nitsche, "Safety limits of cathodal transcranial direct current stimulation in rats," *Clinical Neurophysiology*, vol. 120, no. 6, pp. 1161–1167, 2009.
- [18] M. Jackson, D. Truong, M. Brownlow et al., "Safety parameter considerations of anodal transcranial direct current stimulation in rats," *Brain, Behavior, and Immunity*, vol. 64, pp. 152–161, 2017.
- [19] M. Nitsche, L. Cohen, E. Wassermann et al., "Transcranial direct current stimulation: state of the art 2008," *Brain Stimulation*, vol. 1, no. 3, pp. 206–223, 2008.
- [20] K. Monte-Silva, M. Kuo, D. Liebetanz, W. Paulus, and M. A. Nitsche, "Shaping the optimal repetition interval for cathodal transcranial direct current stimulation (tDCS)," *Journal of Neurophysiology*, vol. 103, no. 4, pp. 1735–1740, 2010.
- [21] P. S. Boggio, R. Ferrucci, S. P. Rigonatti et al., "Effects of transcranial direct current stimulation on working memory in patients with Parkinson's disease," *Journal of the Neurological Sciences*, vol. 249, no. 1, pp. 31–38, 2006.
- [22] M. Horiba, Y. Ueki, I. Nojima et al., "Impaired motor skill acquisition using mirror visual feedback improved by transcranial direct current stimulation (tDCS) in patients with Parkinson's disease," *Frontiers in Neuroscience*, vol. 13, p. 602, 2019.
- [23] F. Fregni, P. S. Boggio, M. A. Nitsche, S. P. Rigonatti, and A. Pascual-Leone, "Cognitive effects of repeated sessions of transcranial direct current stimulation in patients with depression," *Depression and Anxiety*, vol. 23, no. 8, pp. 482–484, 2006.
- [24] F. Fregni, P. Boggio, C. Mansur et al., "Transcranial direct current stimulation of the unaffected hemisphere in stroke patients," *Neuroreport*, vol. 16, no. 14, pp. 1551–1555, 2005.

- [25] F. Hummel, P. Celnik, P. Giraux et al., "Effects of non-invasive cortical stimulation on skilled motor function in chronic stroke," *Brain*, vol. 128, no. 3, pp. 490–499, 2005.
- [26] M. C. W. English, E. S. Kitching, M. T. Maybery, and T. A. W. Visser, "Modulating attentional biases of adults with autistic traits using transcranial direct current stimulation: a pilot study," *Autism Research*, vol. 11, no. 2, pp. 385–390, 2018.
- [27] J. Kang, E. Cai, J. Han et al., "Transcranial direct current stimulation (tDCS) can modulate EEG complexity of children with autism spectrum disorder," *Frontiers in Neuroscience*, vol. 12, p. 201, 2018.
- [28] L. Marshall, M. Mölle, M. Hallschmid, and J. Born, "Transcranial direct current stimulation during sleep improves declarative memory," *Journal of Neuroscience*, vol. 24, no. 44, pp. 9985–9992, 2004.
- [29] J. C. de Souza Custódio, C. W. Martins, M. D. M. V. Lugon, F. Fregni, and E. M. Nakamura-Palacios, "Epidural direct current stimulation over the left medial prefrontal cortex facilitates spatial working memory performance in rats," *Brain Stimulation*, vol. 6, no. 3, pp. 261–269, 2013.
- [30] X. Yu, Y. Li, H. Wen, Y. Zhang, and X. Tian, "Intensity-dependent effects of repetitive anodal transcranial direct current stimulation on learning and memory in a rat model of Alzheimer's disease," *Neurobiology of Learning and Memory*, vol. 123, pp. 168–178, 2015.
- [31] K. J. Yoon, Y. T. Lee, S. W. Chae, C. R. Park, and D. Y. Kim, "Effects of anodal transcranial direct current stimulation (tDCS) on behavioral and spatial memory during the early stage of traumatic brain injury in the rats," *Journal of the Neurological Sciences*, vol. 362, pp. 314–320, 2016.
- [32] T. Kamida, S. Kong, N. Eshima, T. Abe, M. Fujiki, and H. Kobayashi, "Transcranial direct current stimulation decreases convulsions and spatial memory deficits following pilocarpine-induced status epilepticus in immature rats," *Behavioural Brain Research*, vol. 217, no. 1, pp. 99–103, 2011.
- [33] D. T. Leffa, A. de Souza, V. L. Scarabelot et al., "Transcranial direct current stimulation improves short-term memory in an animal model of attention-deficit/hyperactivity disorder," *European Neuropsychopharmacology*, vol. 26, no. 2, pp. 368–377, 2016.
- [34] D. Moher, A. Liberati, J. Tetzlaff, D. G. Altman, and The PRISMA Group, "Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement," *PLoS Medicine*, vol. 6, no. 7, article e1000097, 2009.
- [35] M. R. Krause, T. P. Zanos, B. A. Csorba et al., "Transcranial direct current stimulation facilitates associative learning and alters functional connectivity in the primate brain," *Current Biology*, vol. 27, no. 20, pp. 3086–3096.e3, 2017.
- [36] F. Manteghi, M. Nasehi, and M. Zarrindast, "Precondition of right frontal region with anodal tDCS can restore the fear memory impairment induced by ACPA in male mice," *EXCLI Journal*, vol. 16, pp. 1–13, 2017.
- [37] M. Nasehi, R. Soltanpour, M. Ebrahimi-Ghiri, S. Zarrabian, and M. R. Zarrindast, "Interference effects of transcranial direct current stimulation over the right frontal cortex and adrenergic system on conditioned fear," *Psychopharmacology*, vol. 234, no. 22, pp. 3407–3416, 2017.
- [38] M. Nasehi, M. Khani-Abyaneh, M. Ebrahimi-Ghiri, and M. R. Zarrindast, "The effect of left frontal transcranial direct-current stimulation on propranolol-induced fear memory acquisition and consolidation deficits," *Behavioural Brain Research*, vol. 331, pp. 76–83, 2017.
- [39] S. Abbasi, M. Nasehi, H. R. S. Lichaei, and M. R. Zarrindast, "Effects of left prefrontal transcranial direct current stimulation on the acquisition of contextual and cued fear memory," *Iranian Journal of Basic Medical Sciences*, vol. 20, no. 6, pp. 623–630, 2017.
- [40] M. Podda, S. Cocco, A. Mastrodonato et al., "Anodal transcranial direct current stimulation boosts synaptic plasticity and memory in mice via epigenetic regulation of Bdnf expression," *Scientific Reports*, vol. 6, p. 22180, 2016.
- [41] T. H. Yu, Y. J. Wu, M. E. Chien, and K. S. Hsu, "Transcranial direct current stimulation induces hippocampal metaplasticity mediated by brain-derived neurotrophic factor," *Neuropharmacology*, vol. 144, no. 1, pp. 358–367, 2019.
- [42] C. Dockery, D. Liebetanz, N. Birbaumer, M. Malinowska, and M. J. Wesierska, "Cumulative benefits of frontal transcranial direct current stimulation on visuospatial working memory training and skill learning in rats," *Neurobiology of Learning and Memory*, vol. 96, no. 3, pp. 452–460, 2011.
- [43] S. Yu, S. Park, and K. Sim, "The effect of tDCS on cognition and neurologic recovery of rats with Alzheimer's disease," *Journal of Physical Therapy Science*, vol. 26, no. 2, pp. 247–249, 2014.
- [44] C. H. Chang, H. Y. Lane, and C. H. Lin, "Brain stimulation in Alzheimer's disease," *Frontiers in Psychiatry*, vol. 9, 2018.
- [45] A. Roostaei, G. Vaezi, M. Nasehi, A. Haeri-Rohani, and M. R. Zarrindast, "The involvement of D1 and D2 dopamine receptors in the restoration effect of left frontal anodal, but not cathodal, tDCS on streptozocin-induced amnesia," *Archives of Iranian Medicine*, vol. 22, no. 3, pp. 144–154, 2019.
- [46] Y. Wu, C. Lin, C. Yeh et al., "Repeated transcranial direct current stimulation improves cognitive dysfunction and synaptic plasticity deficit in the prefrontal cortex of streptozotocin-induced diabetic rats," *Brain Stimulation*, vol. 10, no. 6, pp. 1079–1087, 2017.
- [47] D. T. Leffa, B. Bellaver, A. A. Salvi et al., "Transcranial direct current stimulation improves long-term memory deficits in an animal model of attention-deficit/hyperactivity disorder and modulates oxidative and inflammatory parameters," *Brain Stimulation*, vol. 11, no. 4, pp. 743–751, 2018.
- [48] O. Bragina, D. Lara, E. Nemoto, C. W. Shuttleworth, O. V. Semyachkina-Glushkovskaya, and D. E. Bragin, "Increases in microvascular perfusion and tissue oxygenation via vasodilatation after anodal transcranial direct current stimulation in the healthy and traumatized mouse brain," *Advances in Experimental Medicine and Biology*, vol. 1072, pp. 27–31, 2018.
- [49] P. Spagnolo and D. Goldman, "Neuromodulation interventions for addictive disorders: challenges, promise and roadmap for future research," *Brain*, vol. 140, no. 5, pp. 1183–1203, 2017.
- [50] S. Pedron, J. Monnin, E. Haffen, D. Sechter, and V. van Waes, "Repeated transcranial direct current stimulation prevents abnormal behaviors associated with abstinence from chronic nicotine consumption," *Neuropsychopharmacology*, vol. 39, no. 4, pp. 981–988, 2014.
- [51] S. Pedron, J. Beverley, E. Haffen, P. Andrieu, H. Steiner, and V. van Waes, "Transcranial direct current stimulation produces long-lasting attenuation of cocaine-induced behavioral responses and gene regulation in corticostriatal circuits," *Addiction Biology*, vol. 22, no. 5, pp. 1267–1278, 2017.
- [52] T. Kamida, S. Kong, N. Eshima, and M. Fujiki, "Cathodal transcranial direct current stimulation affects seizures and

- cognition in fully amygdala-kindled rats,” *Neurological Research*, vol. 35, no. 6, pp. 602–607, 2013.
- [53] M. Zobeiri and G. van Luijckelaar, “Noninvasive transcranial direct current stimulation in a genetic absence model,” *Epilepsy & Behavior*, vol. 26, no. 1, pp. 42–50, 2013.
- [54] S. C. Dhamne, D. Ekstein, Z. Zhuo et al., “Acute seizure suppression by transcranial direct current stimulation in rats,” *Annals of Clinical and Translational Neurology*, vol. 2, no. 8, pp. 843–856, 2015.
- [55] J. G. Rohan, K. A. Carhuatanta, S. M. McInturf, M. K. Miklasevich, and R. Jankord, “Modulating hippocampal plasticity with in vivo brain stimulation,” *The Journal of Neuroscience*, vol. 35, no. 37, pp. 12824–12832, 2015.
- [56] E. Gondard, M. Soto-Montenegro, A. Cassol, A. M. Lozano, and C. Hamani, “Transcranial direct current stimulation does not improve memory deficits or alter pathological hallmarks in a rodent model of Alzheimer’s disease,” *Journal of Psychiatric Research*, vol. 114, pp. 93–98, 2019.
- [57] K. J. Yoon, B. M. Oh, and D. Y. Kim, “Functional improvement and neuroplastic effects of anodal transcranial direct current stimulation (tDCS) delivered 1 day vs. 1 week after cerebral ischemia in rats,” *Brain Research*, vol. 1452, pp. 61–72, 2012.
- [58] D. Hebb, *The Organization of Behavior*, Wiley, New York, 1949.
- [59] M. Rueger, M. Keuters, M. Walberer et al., “Multi-session transcranial direct current stimulation (tDCS) elicits inflammatory and regenerative processes in the rat brain,” *PLoS One*, vol. 7, no. 8, article e43776, 2012.
- [60] L. Peruzzotti-Jametti, M. Cambiaghi, M. Bacigaluppi et al., “Safety and efficacy of transcranial direct current stimulation in acute experimental ischemic stroke,” *Stroke*, vol. 44, no. 11, pp. 3166–3174, 2013.
- [61] C. McCaig, L. Sangster, and R. Stewart, “Neurotrophins enhance electric field-directed growth cone guidance and directed nerve branching,” *Developmental Dynamics*, vol. 217, no. 3, pp. 299–308, 2000.
- [62] A. Pikhovych, N. P. Stolberg, L. Jessica Flitsch et al., “Transcranial direct current stimulation modulates neurogenesis and microglia activation in the mouse brain,” *Stem Cells International*, vol. 2016, 9 pages, 2016.
- [63] J. Márquez-Ruiz, R. Leal-Campanario, R. Sánchez-Campusano et al., “Transcranial direct-current stimulation modulates synaptic mechanisms involved in associative learning in behaving rabbits,” *Proceedings of the National Academy of Sciences of the United States of America*, vol. 109, no. 17, pp. 6710–6715, 2012.
- [64] B. Krause, J. Márquez-Ruiz, and R. C. Kadosh, “The effect of transcranial direct current stimulation: a role for cortical excitation/inhibition balance?,” *Frontiers in Human Neuroscience*, vol. 7, no. 602, 2013.
- [65] D. Antonenko, F. Schubert, F. Bohm et al., “tDCS-induced modulation of GABA levels and resting-state functional connectivity in older adults,” *The Journal of Neuroscience*, vol. 37, no. 15, pp. 4065–4073, 2017.
- [66] C. Stagg and M. Nitsche, “Physiological basis of transcranial direct current stimulation,” *The Neuroscientist*, vol. 17, no. 1, pp. 37–53, 2011.
- [67] C. J. Stagg, V. Bachtiar, U. Amadi et al., “Local GABA concentration is related to network-level resting functional connectivity,” *eLife*, vol. 3, no. 3, pp. 1–9, 2014.
- [68] B. Fritsch, J. Reis, K. Martinowich et al., “Direct current stimulation promotes BDNF-dependent synaptic plasticity: potential implications for motor learning,” *Neuron*, vol. 66, no. 2, pp. 198–204, 2010.
- [69] F. Ranieri, M. Podda, E. Riccardi et al., “Modulation of LTP at rat hippocampal CA3-CA1 synapses by direct current stimulation,” *Journal of Neurophysiology*, vol. 107, no. 7, pp. 1868–1880, 2012.
- [70] G. Kronberg, M. Bridi, T. Abel, M. Bikson, and L. C. Parra, “Direct current stimulation modulates LTP and LTD: activity dependence and dendritic effects,” *Brain Stimulation*, vol. 10, no. 1, pp. 51–58, 2017.
- [71] Y. Sun, J. Lipton, L. Boyle et al., “Direct current stimulation induces mGluR5-dependent neocortical plasticity,” *Annals of Neurology*, vol. 80, no. 2, pp. 233–246, 2016.
- [72] C. Lüscher and R. Malenka, “NMDA receptor-dependent long-term potentiation and long-term depression (LTP/LTD),” *Cold Spring Harbor Perspectives in Biology*, vol. 4, no. 6, p. a005710, 2012.
- [73] J. Stafford, M. Brownlow, A. Qualley, and R. Jankord, “AMPA receptor translocation and phosphorylation are induced by transcranial direct current stimulation in rats,” *Neurobiology of Learning and Memory*, vol. 150, pp. 36–41, 2018.
- [74] S. Perret, A. Cantereau, J. Audin, B. Dufy, and D. Georgescauld, “Interplay between  $\text{Ca}^{2+}$  release and  $\text{Ca}^{2+}$  influx underlies localized hyperpolarization-induced  $[\text{Ca}^{2+}]_i$  waves in prostatic cells,” *Cell Calcium*, vol. 25, no. 4, pp. 297–311, 1999.
- [75] H. Monai, M. Ohkura, M. Tanaka et al., “Calcium imaging reveals glial involvement in transcranial direct current stimulation-induced plasticity in mouse brain,” *Nature Communications*, vol. 7, p. 11100, 2016.
- [76] T. Mishima, T. Nagai, K. Yahagi et al., “Transcranial direct current stimulation (tDCS) induces adrenergic receptor-dependent microglial morphological changes in mice,” *eNeuro*, vol. 6, no. 5, pp. 1–12, 2019.
- [77] P. Opazo, S. Labrecque, C. Tigaret et al., “CaMKII triggers the diffusional trapping of surface AMPARs through phosphorylation of stargazin,” *Neuron*, vol. 67, no. 2, pp. 239–252, 2010.
- [78] M. S. Kim, H. Koo, S. W. Han et al., “Repeated anodal transcranial direct current stimulation induces neural plasticity-associated gene expression in the rat cortex and hippocampus,” *Restorative Neurology and Neuroscience*, vol. 35, no. 2, pp. 137–146, 2017.
- [79] G. Leal, C. Bramham, and C. Duarte, “BDNF and hippocampal synaptic plasticity,” *Vitamins and Hormones*, vol. 104, pp. 153–195, 2017.
- [80] C. McCaig, A. Rajnicek, B. Song, and M. Zhao, “Controlling cell behavior electrically: current views and future potential,” *Physiological Reviews*, vol. 85, no. 3, pp. 943–978, 2005.
- [81] L. Minichiello, “TrkB signalling pathways in LTP and learning,” *Nature Reviews Neuroscience*, vol. 10, no. 12, pp. 850–860, 2009.
- [82] C. M. Alberini, “Transcription factors in long-term memory and synaptic plasticity,” *Physiological Reviews*, vol. 89, no. 1, pp. 121–145, 2009.
- [83] S. T. Carmichael, I. Archibeque, L. Luke, T. Nolan, J. Momiy, and S. Li, “Growth-associated gene expression after stroke: evidence for a growth-promoting region in peri-infarct

- cortex," *Experimental Neurology*, vol. 193, no. 2, pp. 291–311, 2005.
- [84] Y. Li, N. Jiang, C. Powers, and M. Chopp, "Neuronal damage and plasticity identified by microtubule-associated protein 2, growth-associated protein 43, and cyclin D1 immunoreactivity after focal cerebral ischemia in rats," *Stroke*, vol. 29, no. 9, pp. 1972–1981, 1998.
- [85] T. Jiang, R. X. Xu, A. W. Zhang et al., "Effects of transcranial direct current stimulation on hemichannel pannexin-1 and neural plasticity in rat model of cerebral infarction," *Neurosciences*, vol. 226, pp. 421–426, 2012.
- [86] P. Bargiotas, H. Monyer, and M. Schwaninger, "Hemichannels in cerebral ischemia," *Current Molecular Medicine*, vol. 9, no. 2, pp. 186–194, 2009.
- [87] L. Dimov, A. Franciosi, A. Campos, A. R. Brunoni, and R. L. Pagano, "Top-down effect of direct current stimulation on the nociceptive response of rats," *PLoS One*, vol. 11, no. 4, article e0153506, 2016.
- [88] M. Rioult-Pedotti, D. Friedman, and J. Donoghue, "Learning-induced LTP in neocortex," *Science*, vol. 290, no. 5491, pp. 533–536, 2000.
- [89] R. Malenka and M. Bear, "LTP and LTD: an embarrassment of riches," *Neuron*, vol. 44, no. 1, pp. 5–21, 2004.
- [90] J. Lisman, K. Cooper, M. Sehgal, and A. J. Silva, "Memory formation depends on both synapse-specific modifications of synaptic strength and cell-specific increases in excitability," *Nature Neuroscience*, vol. 21, no. 3, pp. 309–314, 2018.
- [91] M. Legrand, R. Troubat, B. Brizard, A. M. le Guisquet, C. Belzung, and W. el-Hage, "Prefrontal cortex rTMS reverses behavioral impairments and differentially activates c-Fos in a mouse model of post-traumatic stress disorder," *Brain Stimulation*, vol. 12, no. 1, pp. 87–95, 2019.
- [92] V. Moliadze, E. Lyzhko, T. Schmanke, S. Andreas, C. M. Freitag, and M. Siniatchkin, "1 mA cathodal tDCS shows excitatory effects in children and adolescents: insights from TMS evoked N100 potential," *Brain Research Bulletin*, vol. 140, pp. 43–51, 2018.
- [93] F. Fisicaro, G. Lanza, A. A. Grasso et al., "Repetitive transcranial magnetic stimulation in stroke rehabilitation: review of the current evidence and pitfalls," *Therapeutic Advances in Neurological Disorders*, vol. 12, p. 1756286419878317, 2019.
- [94] M. Cantone, A. Bramanti, G. Lanza et al., "Cortical plasticity in depression," *ASN Neuro*, vol. 9, no. 3, 2017.
- [95] U. Palm, F. M. Segmiller, A. N. Epple et al., "Transcranial direct current stimulation in children and adolescents: a comprehensive review," *Journal of Neural Transmission (Vienna)*, vol. 123, no. 10, pp. 1219–1234, 2016.
- [96] X. Leinekugel, I. Khalilov, H. McLean et al., "GABA is the principal fast-acting excitatory transmitter in the neonatal brain," *Advances in Neurology*, vol. 79, pp. 189–201, 1999.
- [97] C. W. Martins, L. C. de Melo Rodrigues, M. A. Nitsche, and E. M. Nakamura-Palacios, "AMPA receptors are involved in prefrontal direct current stimulation effects on long-term working memory and GAP-43 expression," *Behavioural Brain Research*, vol. 362, pp. 208–212, 2019.
- [98] V. Nekhendzy, C. Fender, M. Davies et al., "The antinociceptive effect of transcranial electrostimulation with combined direct and alternating current in freely moving rats," *Anesthesia and Analgesia*, vol. 98, no. 3, pp. 730–7, table of contents, 2004.
- [99] N. O. B. Taib and M. Manto, "Trains of transcranial direct current stimulation antagonize motor cortex hypoexcitability induced by acute hemispherectomy," *Journal of Neurosurgery*, vol. 111, no. 4, pp. 796–806, 2009.
- [100] S. Kim, B. Kim, Y. Ko, M. S. Bang, M. H. Kim, and T. R. Han, "Functional and histologic changes after repeated transcranial direct current stimulation in rat stroke model," *Journal of Korean Medical Science*, vol. 25, no. 10, pp. 1499–1505, 2010.
- [101] G. Laste, W. Caumo, L. Adachi et al., "After-effects of consecutive sessions of transcranial direct current stimulation (tDCS) in a rat model of chronic inflammation," *Experimental Brain Research*, vol. 221, no. 1, pp. 75–83, 2012.
- [102] L. N. S. Adachi, W. Caumo, G. Laste et al., "Reversal of chronic stress-induced pain by transcranial direct current stimulation (tDCS) in an animal model," *Brain Research*, vol. 1489, pp. 17–26, 2012.
- [103] F. Notturmo, M. Pace, F. Zappasodi, E. Cam, C. L. Bassetti, and A. Uncini, "Neuroprotective effect of cathodal transcranial direct current stimulation in a rat stroke model," *Journal of the Neurological Sciences*, vol. 342, no. 1-2, pp. 146–151, 2014.
- [104] C. Lu, Y. Wei, R. Hu, Y. Wang, K. Li, and X. Li, "Transcranial Direct Current Stimulation Ameliorates Behavioral Deficits and Reduces Oxidative Stress in 1-Methyl-4-Phenyl-1,2,3,6-Tetrahydropyridine-Induced Mouse Model of Parkinson's Disease," *Neuromodulation*, vol. 18, no. 6, pp. 442–447, 2015.
- [105] L. N. S. Adachi, A. S. Quevedo, A. de Souza et al., "Exogenously induced brain activation regulates neuronal activity by top-down modulation: conceptualized model for electrical brain stimulation," *Experimental Brain Research*, vol. 233, no. 5, pp. 1377–1389, 2015.
- [106] Y.-H. Liu, L.-D. Liao, S. J. Chan, A. Bandla, and N. V. Thakor, "An integrated neuroprotective intervention for brain ischemia validated by ECoG-fPAM," in *2016 38th Annual International Conference of the IEEE Engineering in Medicine and Biology Society (EMBC)*, pp. 4009–4012, Orlando, FL, USA, Aug. 2016.
- [107] R. Braun, R. Klein, H. Walter et al., "Transcranial direct current stimulation accelerates recovery of function, induces neurogenesis and recruits oligodendrocyte precursors in a rat model of stroke," *Experimental Neurology*, vol. 279, pp. 127–136, 2016.
- [108] S. G. Cioato, L. F. Medeiros, P. R. M. Filho et al., "Long-lasting effect of transcranial direct current stimulation in the reversal of hyperalgesia and cytokine alterations induced by the neuropathic pain model," *Brain Stimulation*, vol. 9, no. 2, pp. 209–217, 2016.
- [109] P. R. M. Filho, R. Vercelino, S. G. Cioato et al., "Transcranial direct current stimulation (tDCS) reverts behavioral alterations and brainstem BDNF level increase induced by neuropathic pain model: Long-lasting effect," *Progress in Neuro-Psychopharmacology & Biological Psychiatry*, vol. 64, pp. 44–51, 2016.
- [110] S. F. da Silva Moreira, L. F. Medeiros, A. de Souza et al., "Transcranial direct current stimulation (tDCS) neuromodulatory effects on mechanical hyperalgesia and cortical BDNF levels in ovariectomized rats," *Life Sciences*, vol. 145, pp. 233–239, 2016.
- [111] Y. Liu, S. Chan, H. Pan et al., "Integrated treatment modality of cathodal-transcranial direct current stimulation with

- peripheral sensory stimulation affords neuroprotection in a rat stroke model,” *Neurophotonic*, vol. 4, no. 4, article 045002, 2017.
- [112] C. Winkler, J. Reis, N. Hoffmann et al., “Anodal transcranial direct current stimulation enhances survival and integration of dopaminergic cell transplants in a rat Parkinson model,” *eNeuro*, vol. 4, no. 5, pp. 1–11, 2017.
- [113] A. Souza, D. F. Martins, L. F. Medeiros et al., “Neurobiological mechanisms of antiallodynic effect of transcranial direct current stimulation (tDCS) in a mice model of neuropathic pain,” *Brain Research*, vol. 1682, pp. 14–23, 2018.
- [114] F. Paciello, M. V. Podda, R. Rolesi et al., “Anodal transcranial direct current stimulation affects auditory cortex plasticity in normal-hearing and noise-exposed rats,” *Brain Stimulation*, vol. 11, no. 5, pp. 1008–1023, 2018.
- [115] F. Fregni, I. Macedo, L. Spezia-Adachi et al., “Transcranial direct current stimulation (tDCS) prevents chronic stress-induced hyperalgesia in rats,” *Brain Stimulation*, vol. 11, no. 2, pp. 299–301, 2018.
- [116] S. Lee, J. Youn, W. Jang, and H. O. Yang, “Neuroprotective effect of anodal transcranial direct current stimulation on 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP)-induced neurotoxicity in mice through modulating mitochondrial dynamics,” *Neurochemistry International*, vol. 129, p. 104491, 2019.
- [117] E. M. M. Callai, V. L. Scarabelot, L. Fernandes Medeiros et al., “Transcranial direct current stimulation (tDCS) and trigeminal pain: a preclinical study,” *Oral Diseases*, vol. 25, no. 3, pp. 888–897, 2019.
- [118] V. Scarabelot, C. de Oliveira, L. Medeiros et al., “Transcranial direct-current stimulation reduces nociceptive behaviour in an orofacial pain model,” *Journal of Oral Rehabilitation*, vol. 46, no. 1, pp. 40–50, 2019.