



Pervin Spaces and Frith Frames: Bitopological Aspects and Completion

Célia Borlido¹ · Anna Laura Suarez²

Received: 6 March 2023 / Accepted: 1 September 2023 / Published online: 30 September 2023
© The Author(s), under exclusive licence to Springer Nature B.V. 2023

Abstract

A Pervin space is a set equipped with a bounded sublattice of its powerset, while its pointfree version, called Frith frame, consists of a frame equipped with a generating bounded sublattice. It is known that the dual adjunction between topological spaces and frames extends to a dual adjunction between Pervin spaces and Frith frames, and that the latter may be seen as representatives of certain quasi-uniform structures. As such, they have an underlying bitopological structure and inherit a natural notion of completion. In this paper we start by exploring the bitopological nature of Pervin spaces and of Frith frames, proving some categorical equivalences involving zero-dimensional structures. We then provide a conceptual proof of a duality between the categories of T_0 complete Pervin spaces and of complete Frith frames. This enables us to interpret several Stone-type dualities as a restriction of the dual adjunction between Pervin spaces and Frith frames along full subcategory embeddings. Finally, we provide analogues of Banaschewski and Pultr's characterizations of sober and T_D topological spaces in the setting of Pervin spaces and of Frith frames, highlighting the parallelism between the two notions.

Keywords Lattice · Pervin space · Distributive lattice · Quasi-uniform space

1 Introduction

The category of Pervin spaces is introduced in [11, 19] as an isomorph of the category of transitive and totally bounded quasi-uniform spaces. Its pointfree analogue, whose objects are named Frith frames, was later defined in [7]. In this setting, we have a full embedding of

Communicated by Matias Menni.

✉ Anna Laura Suarez
annalaurasuares993@gmail.com

Célia Borlido
cborlido@mat.uc.pt

¹ Centre for Mathematics of the University of Coimbra (CMUC), 3001-501 Coimbra, Portugal

² Laboratoire J. A. Dieudonné UMR CNRS 7351, Université Nice Sophia Antipolis, 06108 Nice Cedex 02, France

the category of Frith frames in that of transitive and totally bounded quasi-uniform frames, which is a coreflection but not an equivalence. It is also shown in [7] that the classical dual adjunction between topological spaces and frames naturally extends to a dual adjunction between Pervin spaces and Frith frames. In fact, this is what justifies calling Frith frames the pointfree version of Pervin spaces.

Since both Pervin spaces and Frith frames may be seen as quasi-uniform structures, they come equipped with an underlying bitopological structure as well [10, Chapter 3]. The study of such bitopological structure is the main content of Sect. 3. In Sect. 3.1, we start by assigning a bitopological space to each Pervin space, and show that this is a functorial assignment with a left adjoint. When studying the categorical equivalence induced by such adjunction, *strong exactness* (for Pervin spaces) and *zero-dimensionality* (for bitopological spaces) appear as crucial concepts. More precisely, we show that the categories of the so-called *strongly exact Pervin spaces* and of *zero-dimensional bitopological spaces* are equivalent. We then consider the pointfree version of these results and show that *strongly exact Frith frames* are a full coreflective subcategory of the category of *zero-dimensional biframes*, leaving as an open problem to describe the underlying categorical equivalence. Noting that topological spaces and frames may be seen as bitopological spaces and biframes, respectively, in Sect. 3.2, we specialize the results of the previous section in the monotopological setting. In particular, we show that the categories of *zero-dimensional topological spaces* and of *strongly exact symmetric Pervin spaces* are equivalent, and so are those of *zero-dimensional frames* and of *strongly exact symmetric Frith frames*.

As representatives of quasi-uniform structures, Pervin spaces and Frith frames also inherit natural notions of *completeness*. It is observed in [19] that T_0 and complete Pervin spaces can be identified with spectral spaces, while in [7] it is shown that complete Frith frames can be identified with bounded distributive lattices. In particular, thanks to Stone duality for bounded distributive lattices, it follows that the categories of T_0 and complete Pervin spaces and of complete Frith frames are dual to each other. In Sect. 4, we provide a direct and conceptual proof of this duality, which is based on a characterization of *complete Pervin spaces* and of *complete Frith frames*, and does not invoke Stone duality. On the other hand, since the categories of T_0 complete Pervin spaces and of complete Frith frames are full subcategories of the categories of Pervin spaces and of Frith frames, respectively, we may then see Stone duality as a restriction of the Pervin-Frith dual adjunction along *full* subcategory embeddings (unlike what happens when seeing it as a restriction of the dual adjunction between topological spaces and frames). In Sect. 5, we exhibit several Stone-type dualities as suitable restrictions of the dual adjunction along *full* subcategory embeddings. Section 5.1 is devoted to the already mentioned Stone duality, Sect. 5.2 to Priestley duality, and Sect. 5.3 to bitopological duality. In Sect. 5.4, we provide the global picture of the results thus obtained. It is our concern in Sects. 4 and 5 to point out where the assumption of the *Prime Ideal Theorem* is needed.

Finally, in Sect. 6, starting from Banaschewski and Pultr's characterizations of sober and T_D topological spaces [5, Proposition 4.3], which highlights the parallelism between the two notions, we state and prove analogous results for Pervin spaces, where *sober* is replaced by *complete* and T_D by its equivalent for Pervin spaces (the latter notion having been introduced in [7, Section 4.5]). When looking for a pointfree version of such results, we are naturally led to consider the notion of *locale-based Frith frame*, which will replace T_D Pervin space in our statement.

We readily warn the reader that, although we implicitly have in mind the quasi-uniform interpretation of Pervin spaces and of Frith frames (namely, when considering their bitopological nature and the property of being complete), we will avoid mentioning quasi-uniformities throughout the paper. This reduces the amount of background required from the reader, leav-

ing the details of existing connections for those already familiar with quasi-uniformities. For a detailed study of Pervin spaces, Frith frames, and corresponding quasi-uniform structures, we refer to [7, 11, 19].

2 Preliminaries

The material in this section is presented mostly to set up the notation. We assume the reader to be familiar with frame and locale theory.

2.1 Basic Notation

The content of this section may be found in [13, 18].

A *frame* is a complete lattice L such that for every $a \in L$ and $\{b_i\}_{i \in I} \subseteq L$ the following distributivity law holds:

$$a \wedge \bigvee_{i \in I} b_i = \bigvee_{i \in I} (a \wedge b_i).$$

A frame L is always a complete Heyting algebra, with the *Heyting implication* given by

$$a \rightarrow b = \bigvee \{x \in L \mid x \wedge a \leq b\},$$

for every $a, b \in L$. The element $a \rightarrow 0$, called the *pseudocomplement* of a , will be denoted by a^* . When $a \vee a^* = 1$, we say that a is *complemented* and a^* is the *complement* of a . A *frame homomorphism* is a map $h : L \rightarrow M$ that preserves finite meets and arbitrary joins. We will denote by **Frm** the category of frames and frame homomorphisms. A frame homomorphism $h : L \rightarrow M$ is *dense* provided $h(a) = 0$ implies $a = 0$. In general, a frame homomorphism need not preserve the Heyting implication. We have however the following:

$$h(a \rightarrow b) \leq h(a) \rightarrow h(b), \tag{1}$$

for every $a, b \in L$. Moreover, since every frame homomorphism h preserves arbitrary joins, it has a right adjoint h_* , and the equality

$$a \rightarrow h_*(x) = h_*(h(a) \rightarrow x) \tag{2}$$

holds, for every $a \in L$ and $x \in M$. This is usually called the *Frobenius identity*.

We say that an element $a \in L$ is *compact* if whenever $a \leq \bigvee_{i \in I} a_i$ there exists a finite subset $I' \subseteq I$ such that $a \leq \bigvee_{i \in I'} a_i$. A frame L is *compact* if its top element is compact. If the set of compact elements of L is closed under finite meets and join-generates L , then we say that L is *coherent*. A frame L is *zero-dimensional* if it is join-generated by its sublattice of complemented elements.


The opposite category of **Frm**, is usually denoted by **Loc**. Its objects are called *locales*, and morphisms $h : L \rightarrow M$ are the right adjoints of the corresponding frame maps. Locales will only be mentioned in Sect. 6. For our purposes, the notion of *sublocale* will be enough. A *sublocale* of L is a subset $K \subseteq L$ that is closed under arbitrary joins and contains every element of the form $a \rightarrow x$, with $a \in L$ and $x \in K$. A sublocale K is itself a frame, but not a *subframe* of L , as joins may be computed differently. Every sublocale is uniquely determined by a frame quotient $q : L \twoheadrightarrow K$ whose right adjoint is the localic embedding $K \hookrightarrow L$. In particular, $q(x) = x$, whenever $x \in K$.

Since sublocales correspond to frame quotients, these may also be defined via *frame congruences*, which will be widely used throughout the paper. The set \mathcal{CL} of all frame congruences on L is itself a frame when ordered by inclusion. For every element $a \in L$, we may define two congruences:

$$\nabla_a := \{(x, y) \in L \times L \mid a \vee x = a \vee y\} \quad \text{and} \quad \Delta_a := \{(x, y) \in L \times L \mid a \wedge x = a \wedge y\}.$$

Congruences of the form ∇_a are called *closed*, while those of the form Δ_a are *open*. Open and closed congruences suffice to generate \mathcal{CL} , as a frame. The functions $\nabla : a \mapsto \nabla_a$ and $\Delta : a \mapsto \Delta_a$ from L to \mathcal{CL} are, respectively, a frame embedding and an injection that turns finite meets into finite joins and arbitrary joins to arbitrary meets. Given a subset $S \subseteq L$, we denote by ∇S and by ΔS the subframes of \mathcal{CL} generated by $\{\nabla_s \mid s \in S\}$ and by $\{\Delta_s \mid s \in S\}$, respectively. The subframe of \mathcal{CL} generated by $\nabla L \cup \Delta S$ will be denoted by $\mathcal{C}_S L$. The following generalizes the well-known universal property of the congruence frame.

Proposition 2.1 ([24, Theorem 16.2]) *For every frame L and subset $S \subseteq L$, the frame $\mathcal{C}_S L$ has the following universal property: whenever $h : L \rightarrow M$ is a frame map such that $h(s)$ is complemented for all $s \in S$, there is a unique frame homomorphism $\tilde{h} : \mathcal{C}_S L \rightarrow M$ making the following diagram commute.*



Finally, we have an idempotent adjunction $\Omega : \mathbf{Top} \rightleftarrows \mathbf{Loc} : \mathbf{pt}$ between the category **Top** of topological spaces and continuous functions and the category of locales. Since, as already mentioned, we will mostly work with frames, we will treat the category **Loc** as that opposite to **Frm**. Given a topological space (X, τ) (or simply X if no confusion arises), $\Omega(X)$ is the frame $\Omega(X)$ consisting of the open subsets of X (ordered by \subseteq) and, for a continuous function $f : X \rightarrow Y$, $\Omega(f) = f^{-1}$ is the preimage frame homomorphism. In the other direction, given a frame L , $\mathbf{pt}(L)$ is the set $\mathbf{pt}(L)$ of *points* of L (here seen as frame homomorphisms $p : L \rightarrow \mathbf{2}$) equipped with the topology $\widehat{L} := \{\widehat{a} \mid a \in L\}$, where $\widehat{a} := \{p \in \mathbf{pt}(L) \mid p(a) = 1\}$, and given a frame homomorphism $h : L \rightarrow M$, $\mathbf{pt}(h)$ maps $p \in \mathbf{pt}(M)$ to $p \circ h \in \mathbf{pt}(L)$. The fixpoints of $\Omega \dashv \mathbf{pt}$ are the so-called *sober spaces* and *spatial frames*, respectively.

2.2 Bitopological Spaces and Biframes

We refer to [21] and to [4, 22] for further reading on bitopological spaces and on biframes, respectively.

A *bitopological space*, or *bispace*, is a triple $\mathcal{X} = (X, \tau_+, \tau_-)$, where X is a set and τ_+ and τ_- two topologies on that set. Morphisms between bitopological spaces are functions between their underlying sets that are continuous with respect to both topologies. We denote by **BiTop** the category thus obtained. The topology $\tau_+ \vee \tau_-$ on X generated by $\tau_+ \cup \tau_-$ is called the *patch topology*. We call τ_+ the *positive topology*, and τ_- the *negative one*. Accordingly, elements of τ_+ are called *positive opens*, and a positive open whose complement is a negative one is called a *positive clopen*. The collection of positive clopen subsets of X will be denoted by $\mathbf{Clop}_+(\mathcal{X})$. Negative (c)opens and $\mathbf{Clop}_-(\mathcal{X})$ are defined similarly, in the obvious way.

We say that a bispaces is T_0 (respectively, *compact*) if its patch topology is T_0 (respectively, compact). We say that a bispaces (X, τ_+, τ_-) is *zero-dimensional* if every element in τ_+ is a union of positive clopens, and every element in τ_- is a union of negative clopens.¹

A *biframe* is a triple $\mathcal{L} = (L, L_+, L_-)$ such that all three components are frames, together with subframe inclusions $L_+ \subseteq L$ and $L_- \subseteq L$, and such that every element of L is a join of finite meets of elements of $L_+ \cup L_-$. The frame L is called the *main component* of the biframe, while L_+ and L_- are, respectively, the *positive* and *negative* components. Accordingly, elements of L_+ are *positive* and those of L_- are *negative*. We say that an element $a \in L_+$ is *positive bicomplemented* when it is complemented in L with complement in L_- . *Negative bicomplemented* elements are defined similarly. We denote by $B_+(\mathcal{L})$ and by $B_-(\mathcal{L})$ the lattices of positive and negative bicomplemented elements of \mathcal{L} , respectively. A morphism $h : (L, L_+, L_-) \rightarrow (M, M_+, M_-)$ between biframes is a frame homomorphism $h : L \rightarrow M$ such that $h[L_+] \subseteq M_+$ and $h[L_-] \subseteq M_-$. The category of biframes and biframe homomorphisms will be denoted by **BiFrm**. We say that h is *dense* if its underlying frame homomorphism is dense.

We say that a biframe $\mathcal{L} = (L, L_+, L_-)$ is *compact* if its main component L is compact, and that it is *zero-dimensional* if both L_+ and L_- are join-generated by their bicomplemented elements.

Finally, we also have an adjunction $\Omega_b : \mathbf{BiTop} \rightleftharpoons \mathbf{BiFrm}^{op} : \mathbf{pt}_b$ between bitopological spaces and biframes, which extends the classical adjunction between topological spaces and frames (here, we identify a topological space (X, τ) with the bispaces (X, τ, τ) , and a frame L with the biframe (L, L, L)). Given a bitopological space $\mathcal{X} = (X, \tau_+, \tau_-)$, $\Omega_b(\mathcal{X})$ is the biframe $(\tau_+ \vee \tau_-, \tau_+, \tau_-)$, while for a biframe $\mathcal{L} = (L, L_+, L_-)$, $\mathbf{pt}_b(\mathcal{L}) = (\text{pt}(L), \widehat{L}_+, \widehat{L}_-)$, where for $P \in \{L_+, L_-\}$, we denote $\widehat{P} := \{\widehat{a} \mid a \in P\}$. On morphisms, Ω_b and \mathbf{pt}_b are defined as expected.

2.3 The Prime Ideal Theorem

One of the main features of *pointfree topology* is that it often avoids the use of the Axiom of Choice, thereby leading to constructive results. Sometimes, a strictly weaker version known as the *Prime Ideal Theorem* is however needed. It may be stated as follows:

Let D be a bounded distributive lattice. Then, every proper ideal of D may be extended to a prime ideal.

In Sects. 4 and 5, some results are valid only under the assumption of the Prime Ideal Theorem. The following equivalent statements will be useful in there.

Theorem 2.2 *The following statements are equivalent:*

- (a) *The Prime Ideal Theorem holds.*
- (b) *For every set X , every proper filter of $\mathcal{P}(X)$ can be extended to an ultrafilter.*
- (c) *Every frame of the form $\text{Idl}(D)$, where D is a bounded distributive lattice, is spatial.*

Relevant references for the equivalence between these statements are [2, 9, 12, 13].

¹ These notions are not consistent over all the literature. For instance, in [6] our notions of T_0 and compact are named *join T_0* and *join compact*, respectively, while compact is named *pairwise compact* in [21]. Moreover, in [6, 20] our notion of zero-dimensional is named *pairwise zero-dimensional*.

2.4 Pervin Spaces and Frith Frames

A *Pervin space* is a pair (X, \mathcal{S}) where X is a set and \mathcal{S} a sublattice of its powerset. A morphism $f : (X, \mathcal{S}) \rightarrow (Y, \mathcal{T})$ of Pervin spaces is a set function $f : X \rightarrow Y$ such that $f^{-1}(T) \subseteq \mathcal{S}$. Since a topology on a set X is, in particular, a bounded sublattice of its powerset, the category **Top** of topological spaces fully embeds in the category **Pervin** of Pervin spaces.

Proposition 2.3 [7, Section 3] *Let $f : (X, \mathcal{S}) \rightarrow (Y, \mathcal{T})$ be a map of Pervin spaces. Then,*

- (a) *f is an epimorphism if and only if its underlying set map is surjective;*
- (b) *f is an extremal monomorphism if and only if its underlying set map is injective and every element of \mathcal{S} is of the form $f^{-1}(T)$ for some $T \in \mathcal{T}$.*

In particular, f is an isomorphism if and only if its underlying set map is a bijection and $f[\mathcal{S}] = \mathcal{T}$.

Recall that a *join-dense* subset of a complete lattice L is a subset whose closure under (arbitrary) joins is L itself. A *Frith frame* is a pair (L, \mathcal{S}) , where L is a frame and \mathcal{S} is a join-dense bounded sublattice of L . A morphism of Frith frames $h : (L, \mathcal{S}) \rightarrow (M, \mathcal{T})$ is a frame homomorphism between the underlying frames that satisfies $h[\mathcal{S}] \subseteq \mathcal{T}$. The category of Frith frames and their morphisms is denoted **Frith**, and we have a full embedding **Frm** \hookrightarrow **Frith** obtained by identifying a frame L with the Frith frame (L, L) .

Proposition 2.4 ([7, Section 4.4]) *Let $h : (L, \mathcal{S}) \rightarrow (M, \mathcal{T})$ be a homomorphism of Frith frames. Then,*

- (a) *h is a monomorphism if and only if so is its underlying frame homomorphism;*
- (b) *h is an extremal epimorphism if and only if it satisfies $h[\mathcal{S}] = \mathcal{T}$.*

In particular, h is an isomorphism if and only if it is injective and satisfies $h[\mathcal{S}] = \mathcal{T}$.

The classical dual adjunction between topological spaces and frames may then be extended to a dual adjunction $\Omega : \mathbf{Pervin} \rightleftarrows \mathbf{Frith} : \mathbf{pt}$ between Pervin spaces and Frith frames as follows. For a Pervin space (X, \mathcal{S}) , $\Omega(X, \mathcal{S})$ is the Pervin space $(\Omega_{\mathcal{S}}(X), \mathcal{S})$, where $\Omega_{\mathcal{S}}(X)$ denotes the topology on X generated by \mathcal{S} . For a morphism $f : (X, \mathcal{S}) \rightarrow (Y, \mathcal{T})$, $\Omega(f)$ is the preimage map $f^{-1} : (\Omega_{\mathcal{T}}(Y), \mathcal{T}) \rightarrow (\Omega_{\mathcal{S}}(X), \mathcal{S})$. In the other direction, for a Frith frame (L, \mathcal{S}) , $\mathbf{pt}(L, \mathcal{S})$ is the Pervin space $(\mathbf{pt}(L), \widehat{\mathcal{S}})$, where $\widehat{\mathcal{S}} := \{\widehat{s} \mid s \in \mathcal{S}\}$. Finally, given a morphism of Frith frames $h : (L, \mathcal{S}) \rightarrow (M, \mathcal{T})$, $\mathbf{pt}(h)$ maps $p \in \mathbf{pt}(M)$ to $p \circ h \in \mathbf{pt}(L)$.

Theorem 2.5 ([7, Proposition 4.3]) *There is an adjunction $\Omega : \mathbf{Pervin} \rightleftarrows \mathbf{Frith}^{op} : \mathbf{pt}$ with $\Omega \dashv \mathbf{pt}$, whose fixpoints are, respectively, the Pervin spaces (X, \mathcal{S}) such that $(X, \Omega_{\mathcal{S}}(X))$ is sober and the Frith frames (L, \mathcal{S}) such that L is spatial. These will be called, respectively, sober Pervin spaces and spatial Frith frames.*

2.5 Symmetrization

Symmetrization is for Pervin spaces and Frith frames, what *uniformization* is for quasi-uniform spaces and quasi-uniform frames, respectively. This has been considered in [19] for Pervin spaces and in [7] for Frith frames. In particular, it has been shown that the categories of *symmetric* Pervin spaces and of *symmetric* Frith frames are equivalent to the categories of transitive and totally bounded uniform spaces and frames, respectively.

Recall that a Pervin space (X, \mathcal{B}) is *symmetric* if \mathcal{B} is a Boolean algebra, and the full subcategory of **Pervin** determined by the symmetric Pervin spaces will be denoted by **Pervin**_{sym}. A Frith frame (L, B) is *symmetric* if B is a Boolean algebra, and the full subcategory of **Frith** determined by the symmetric Frith frames will be denoted by **Frith**_{sym}.

We define a functor **Sym**_{Perv} : **Pervin** → **Pervin**_{sym} as follows. For an object (X, \mathcal{S}) , we let **Sym**_{Perv} (X, \mathcal{S}) be the Pervin space $(X, \overline{\mathcal{S}})$, where $\overline{\mathcal{S}}$ is the Boolean subalgebra of the powerset $\mathcal{P}(X)$ generated by the elements of \mathcal{S} . On morphisms, we simply map a function to itself.

Proposition 2.6 ([7, Proposition 3.4]) *The functor **Sym**_{Perv} is right adjoint to the embedding **Pervin**_{sym} ↪ **Pervin**.*


The pointfree version of **Sym**_{Perv} is defined as follows. For a Frith frame (L, S) we set **Sym**_{Frith} $(L, S) := (C_S L, \overline{S})$, where \overline{S} denotes the sublattice of $C_S L$ generated by the elements of the form ∇_s together with their complements. For a morphism of Frith frames $h : (L, S) \rightarrow (M, T)$ we set **Sym**_{Frith} $(h) := \overline{h}$, where \overline{h} is the unique extension of h to a frame homomorphism $\overline{h} : C_S L \rightarrow C_T M$ (recall Proposition 2.1).

Proposition 2.7 ([7, Proposition 6.5]) *The functor **Sym**_{Frith} is left adjoint to the embedding **Frith**_{sym} ↪ **Frith**.*

2.6 Forgetful Functors

Both Pervin spaces and Frith frames encode two kinds of structures - topological and lattice-theoretical. It is then natural to consider several forgetful functors on the categories **Pervin** and **Frith**.

Every Pervin space (X, \mathcal{S}) defines a topology $\Omega_{\mathcal{S}}(X)$ on X . This assignment can be extended to a functor **U**_{Pervin} : **Pervin** → **Top** which leaves functions unaltered.² The pointfree version of this functor is the functor **U**_{Frith} : **Frith** → **Frm**, which acts as $(L, S) \mapsto L$ on objects, and which leaves frame maps unaltered. The relationship between **U**_{Pervin} and **U**_{Frith} is depicted in the following commutative diagrams:



Proposition 2.8 ([7]) *The functor **U**_{Pervin} is right adjoint to the embedding **Top** ↪ **Pervin**, and the functor **U**_{Frith} is left adjoint to the embedding **Frm** ↪ **Frith**.*

Given a topological property, we will often say that a Pervin space has that property provided so does its underlying topological space. For instance, a Pervin space (X, \mathcal{S}) is T_0 if $(X, \Omega_{\mathcal{S}}(X))$ is T_0 . The same applies to the notion of *dense* morphism. We say that a morphism $f : (X, \mathcal{S}) \rightarrow (Y, \mathcal{T})$ is *dense* if the image of $\mathbf{U}_{\text{Pervin}}(f) : (X, \Omega_{\mathcal{S}}(X)) \rightarrow (Y, \Omega_{\mathcal{T}}(Y))$ is dense in $(Y, \Omega_{\mathcal{T}}(Y))$, that is, if $f[X]$ intersects every nonempty open subset of $(Y, \Omega_{\mathcal{T}}(Y))$. The following characterization of density for Pervin morphisms is easy to prove. We will use it without further mention.

² Under the identification of Pervin spaces with transitive and totally bounded quasi-uniform spaces, this functor forgets the quasi-uniform structure.

Lemma 2.9 For a map $f : (X, S) \rightarrow (Y, T)$ of Pervin spaces, the following are equivalent.

- (a) The map f is dense;
- (b) $f[X]$ intersects every nonempty element of T ;
- (c) We have $f^{-1}(T) = \emptyset$ implies $T = \emptyset$ for all $T \in \mathcal{T}$.

We may also forget the topological structures, thereby obtaining functors $\mathbf{L}_{\text{Pervin}} : \mathbf{Pervin} \rightarrow \mathbf{DLat}^{op}$ and $\mathbf{L}_{\text{Frith}} : \mathbf{Frith} \rightarrow \mathbf{DLat}$ defined in the expected way.

In [7, Proposition 4.6], we have proved that $\mathbf{L}_{\text{Frith}}$ is right adjoint to the functor $\mathbf{Idl} : \mathbf{DLat} \rightarrow \mathbf{Frith}$ acting on objects as $S \mapsto (\text{Idl}(S), S)$ and assigning to each morphism $f : S \rightarrow T$ its unique extension to a frame morphism $\text{Idl}(S) \rightarrow \text{Idl}(T)$. The component of the counit of this adjunction at a Frith frame (L, S) is the morphism

$$c_{(L,S)} : (\text{Idl}(S), S) \rightarrow (L, S), \quad J \mapsto \bigvee J. \tag{3}$$

We now discuss the point-set version of this result. To define the right adjoint of $\mathbf{L}_{\text{Pervin}}$, we introduce some notation. Given a lattice D , we denote by $\text{pf}(D)$ the set of all prime filters of D , and we consider the map

$$\Phi_D : D \rightarrow \mathcal{P}(\text{pf}(D)), \quad a \mapsto \tilde{a} := \{F \in \text{pf}(D) \mid a \in F\}. \tag{4}$$

We denote as $\tilde{D} := \{\tilde{a} \mid a \in D\}$. It is well-known that Φ_D is a lattice homomorphism and it is an embedding if and only if D is isomorphic to a sublattice of the powerset $\mathcal{P}(X)$ for some set X (see e.g. [9, Chapter 10]). If we further assume that the Prime Ideal Theorem holds, then Φ_D is always an embedding.

We define the functor $\mathbf{pf} : \mathbf{DLat}^{op} \rightarrow \mathbf{Pervin}$ as the one mapping each lattice D to the Pervin space $(\text{pf}(D), \tilde{D})$, and each lattice homomorphism $f : D_1 \rightarrow D_2$ to the preimage map $\mathbf{pf}(f) := f^{-1} : (\text{pf}(D_2), \tilde{D}_2) \rightarrow (\text{pf}(D_1), \tilde{D}_1)$. This is well-defined as preimages of prime filters are prime filters, and $\mathbf{pf}(f)^{-1}(\tilde{a}) = \tilde{f(a)}$ for every $a \in D_1$.

Lemma 2.10 We have an idempotent adjunction $\mathbf{L}_{\text{Pervin}} : \mathbf{Pervin} \rightleftarrows \mathbf{DLat}^{op} : \mathbf{pf}$ with $\mathbf{L}_{\text{Pervin}} \dashv \mathbf{pf}$.

Proof With a routine computation one can show that, given a lattice D , the co-restriction of Φ_D to a map $D \rightarrow \tilde{D}$ is a universal morphism from $\mathbf{L}_{\text{Pervin}}$ to D and thus, the counit of the adjunction at D . To conclude that the adjunction is idempotent, we only have to note that Φ_D is an isomorphism whenever D is a sublattice of some powerset and that is the case for every lattice component of a Pervin space (X, S) . □

We further remark that the component of the unit of the adjunction $\mathbf{L}_{\text{Pervin}} \dashv \mathbf{pf}$ at a Pervin space (X, S) is the neighborhood morphism of Pervin spaces defined by

$$\mathcal{N}_{(X,S)} : (X, S) \rightarrow (\text{pf}(S), \tilde{S}), \quad x \mapsto \{S \in \mathcal{S} \mid x \in S\}. \tag{5}$$

We finish this section by providing an alternative representation of the Pervin space $\mathbf{pf}(D)$, based on the well-known correspondence between prime filters of D and points of $\text{Idl}(D)$.

Lemma 2.11 There is a one-to-one correspondence between prime filters of D and points of $\text{Idl}(D)$ and, under this identification, \tilde{a} corresponds to \hat{a} , for every $a \in D$. In particular, $\mathbf{pf}(D)$ is isomorphic to $\mathbf{pt}(\text{Idl}(D), D)$.

2.7 Open Intersections and Strongly Exact Meets

For a Pervin space (X, \mathcal{S}) we say that an intersection of elements in \mathcal{S} is *open* if it is so in the topological space $(X, \Omega_{\mathcal{S}}(X))$. We will denote by $[\mathcal{S}]_{op}$ the collection of all elements of $\Omega_{\mathcal{S}}(X)$ that are open intersections of elements of \mathcal{S} . It is easy to see that this collection is closed under open intersections and so, $[\mathcal{S}]_{op}$ may be seen as the closure of \mathcal{S} under open intersections.

Let now L be a frame and $P \subseteq L$. The meet $\bigwedge P$ is *strongly exact* if the corresponding intersection of open sublocales (=open pointfree subspaces) is open, or equivalently, if the congruence $\bigvee_{s \in P} \Delta_s$ is open (cf. [1, Section 4.5]). Note that, if $\Delta_a = \bigvee_{s \in P} \Delta_s$ for some $P \subseteq L$, then the meet $\bigwedge P$ is, by definition, strongly exact, and we necessarily have $a = \bigwedge P$. Given a Frith frame (L, \mathcal{S}) , we denote by $[\mathcal{S}]_{se}$ the set of elements of L that may be written as a strongly exact meet of elements of \mathcal{S} . Again, $[\mathcal{S}]_{se}$ can be thought of as the closure of \mathcal{S} under strongly exact meets.

Finally, we say that a Pervin space (X, \mathcal{S}) (respectively, Frith frame (L, \mathcal{S})) is *strongly exact* if the lattice \mathcal{S} (respectively, \mathcal{S}) is closed under open intersections of \mathcal{S} (respectively, strongly exact meets of \mathcal{S}). We denote by **Pervin**_{se} and by **Frith**_{se} the full subcategories of **Pervin** and of **Frith** determined by the strongly exact objects.

The next result implies that the contravariant functor $\Omega : \mathbf{Pervin} \rightarrow \mathbf{Frith}$ restricts and co-restricts to a functor **Pervin**_{se} \rightarrow **Frith**_{se}.

Proposition 2.12 ([1, Proposition 5.3]) *Let (X, τ) be a topological space and $\mathcal{U} \subseteq \Omega(X)$ be a family of open subsets. If $\bigwedge \mathcal{U}$ is a strongly exact meet in the frame $\Omega(X)$, then $\bigcap \mathcal{U}$ is an open subset of X .*

We do not know whether the functor **pt** : **Frith** \rightarrow **Pervin** restricts and co-restricts to a functor **Frith**_{se} \rightarrow **Pervin**_{se}.

3 The Bitopological Point of View

3.1 Strong Exactness and Zero-Dimensionality

A Pervin space (X, \mathcal{S}) defines the bitopological space $(X, \Omega_{\mathcal{S}}(X), \Omega_{\mathcal{S}^c}(X))$, where \mathcal{S}^c denotes the lattice $\{S^c \mid S \in \mathcal{S}\}$ formed by the complements in X of the elements of \mathcal{S} .³ The *Skulafunctor* **Sk**_{Pervin} : **Pervin** \rightarrow **BiTop** is then defined by assigning $(X, \Omega_{\mathcal{S}}(X), \Omega_{\mathcal{S}^c}(X))$ to the Pervin space (X, \mathcal{S}) , and mapping each function to itself.

In the other direction, we may define a functor **Clop**₊ : **BiTop** \rightarrow **Pervin** by assigning to each bitopological space $\mathcal{X} = (X, \tau_+, \tau_-)$ the Pervin space $(X, \text{Clop}_+(\mathcal{X}))$ and keeping morphisms unchanged.

It is easily seen that these are well-defined functors. Let us prove that **Clop**₊ is left adjoint to **Sk**_{Pervin}.

Lemma 3.1 *The functor **Clop**₊ is left adjoint to **Sk**_{Pervin} : **Pervin** \rightarrow **BiTop**.*

Proof Let $\mathcal{X} = (X, \tau_+, \tau_-)$ be a bitopological space. We first observe that

$$\mathbf{Sk}_{\text{Pervin}} \circ \mathbf{Clop}_+(\mathcal{X}) = (X, \Omega_{\text{Clop}_+(\mathcal{X})}(X), \Omega_{\text{Clop}_-(\mathcal{X})}(X)).$$

³ For the interested reader, this is the underlying bitopological space of the quasi-uniform space represented by (X, \mathcal{S}) .

Since the inclusions $\text{Clop}_+(\mathcal{X}) \subseteq \tau_+$ and $\text{Clop}_-(\mathcal{X}) \subseteq \tau_-$ hold, the identity function on X induces a morphism of bitopological spaces $\eta_{\mathcal{X}} : \mathcal{X} \rightarrow \mathbf{Sk}_{\text{Pervin}} \circ \mathbf{Clop}_+(\mathcal{X})$. Let us show that $(\mathbf{Clop}_+(\mathcal{X}), \eta_{\mathcal{X}})$ is a universal morphism from \mathcal{X} to $\mathbf{Sk}_{\text{Pervin}}$. Let (Y, \mathcal{T}) be a Pervin space and $f : \mathcal{X} \rightarrow \mathbf{Sk}_{\text{Pervin}}(Y, \mathcal{T})$ be a morphism of bitopological spaces. Since $f^{-1}(\mathcal{T}) \subseteq \tau_+$ and $f^{-1}(\mathcal{T}^c) \subseteq \tau_-$, the underlying set function of f defines a morphism $g : \mathbf{Clop}_+(\mathcal{X}) \rightarrow (Y, \mathcal{T})$. Clearly, g is the unique morphism satisfying $\mathbf{Sk}_{\text{Pervin}}(g) \circ \eta_{\mathcal{X}} = f$, and this proves our claim. \square

We now describe the equivalence of categories determined by $\mathbf{Clop}_+ \dashv \mathbf{Sk}_{\text{Pervin}}$.

Proposition 3.2 *The fixpoints of the adjunction $\mathbf{Clop}_+ : \mathbf{BiTop} \rightleftarrows \mathbf{Pervin} : \mathbf{Sk}_{\text{Pervin}}$ are, respectively, the zero-dimensional bispaces and the strongly exact Pervin spaces.*

Proof It follows from the proof of Lemma 3.1 that the unit of the adjunction $\mathbf{Clop}_+ \dashv \mathbf{Sk}_{\text{Pervin}}$ at a bispace $\mathcal{X} = (X, \tau_+, \tau_-)$ is the morphism

$$\eta_{\mathcal{X}} : (X, \tau_+, \tau_-) \rightarrow (X, \Omega_{\text{Clop}_+(\mathcal{X})}(X), \Omega_{\text{Clop}_-(\mathcal{X})}(X))$$

defined by the identity map on X . Thus, \mathcal{X} is a fixpoint of the adjunction if and only if $\tau_+ = \Omega_{\text{Clop}_+(\mathcal{X})}(X)$ and $\tau_- = \Omega_{\text{Clop}_-(\mathcal{X})}(X)$, that is, if and only if \mathcal{X} is zero-dimensional.

Let us now exhibit the counit of $\mathbf{Clop}_+ \dashv \mathbf{Sk}_{\text{Pervin}}$. It is easy to see that the positive clopens of $\mathbf{Sk}_{\text{Pervin}}(X, \mathcal{S}) = (X, \Omega_{\mathcal{S}}(X), \Omega_{\mathcal{S}^c}(X))$ are the open intersections of \mathcal{S} . Since $\mathcal{S} \subseteq [\mathcal{S}]_{op}$, the identity on X defines a morphism

$$\varepsilon_{(X, \mathcal{S})} : (X, [\mathcal{S}]_{op}) \rightarrow (X, \mathcal{S})$$

of Pervin spaces, which can be shown to be the counit of the adjunction. In particular, we have that (X, \mathcal{S}) is a fixpoint if and only if $\mathcal{S} = [\mathcal{S}]_{op}$, that is, if and only if (X, \mathcal{S}) is a strongly exact Pervin space. \square

Corollary 3.3 *The categories \mathbf{BiTop}_Z and \mathbf{Pervin}_{se} of zero-dimensional bitopological spaces and of strongly exact Pervin spaces are equivalent.*

We finally remark that, since, for every Pervin space (X, \mathcal{S}) , we have

$$\begin{aligned} \mathbf{Sk}_{\text{Pervin}} \circ \mathbf{Clop}_+ \circ \mathbf{Sk}_{\text{Pervin}}(X, \mathcal{S}) &= \mathbf{Sk}_{\text{Pervin}}(X, [\mathcal{S}]_{op}) = (X, \Omega_{[\mathcal{S}]_{op}}(X), \Omega_{([\mathcal{S}]_{op})^c}(X)) \\ &= \mathbf{Sk}_{\text{Pervin}}(X, \mathcal{S}), \end{aligned}$$

the unit $\eta_{\mathbf{Sk}_{\text{Pervin}}(X, \mathcal{S})}$ is always an isomorphism, and thus, the adjunction $\mathbf{Clop}_+ \dashv \mathbf{Sk}_{\text{Pervin}}$ is idempotent.

Let us now look at the pointfree version of the Skula functor and its left adjoint. For a Frith frame (L, S) we set $\mathbf{Sk}_{\text{Frith}}(L, S) = (C_S L, \nabla L, \Delta S)$.⁴ In order to define $\mathbf{Sk}_{\text{Frith}}$ on morphisms, we first observe that, by Proposition 2.1, every morphism of Frith frames $h : (L, S) \rightarrow (M, T)$ uniquely extends to a frame homomorphism $\bar{h} : C_S L \rightarrow C_T M$ satisfying

$$\bar{h}[\nabla L] = \nabla h[L] \quad \text{and} \quad \bar{h}[\Delta S] = \Delta h[S].$$

Therefore, \bar{h} defines a biframe homomorphism $\bar{h} : \mathbf{Sk}_{\text{Frith}}(L, S) \rightarrow \mathbf{Sk}_{\text{Frith}}(M, T)$ and we may set $\mathbf{Sk}_{\text{Frith}}(h) = \bar{h}$.

In the other direction, we define $\mathbf{B}_+ : \mathbf{BiFrm} \rightarrow \mathbf{Frith}$ as follows. For a biframe $\mathcal{L} = (L, L_+, L_-)$, we set $\mathbf{B}_+(\mathcal{L}) = ((\mathbf{B}_+(\mathcal{L}))_{\text{Frm}}, \mathbf{B}_+(\mathcal{L}))$, where $(\mathbf{B}_+(\mathcal{L}))_{\text{Frm}}$ denotes the

⁴ As for spaces, this is the underlying biframe of the quasi-uniform frame defined by (L, S) .

subframe of L_+ generated by the lattice $\mathbf{B}_+(\mathcal{L})$ of positive bicomplemented elements of \mathcal{L} . In order to define \mathbf{B}_+ on morphisms, notice that, if $h : \mathcal{L} \rightarrow \mathcal{K}$ is a biframe homomorphism then, since positive bicomplemented elements of \mathcal{L} are mapped to positive bicomplemented elements of \mathcal{K} , the suitable restriction and co-restriction of h induces a morphism of Frith frames $\mathbf{B}_+h : \mathbf{B}_+(\mathcal{L}) \rightarrow \mathbf{B}_+(\mathcal{K})$.

Next we will see that, as for Pervin spaces, the functors $\mathbf{Sk}_{\text{Frith}}$ and \mathbf{B}_+ define an adjunction between the categories of Frith frames and of biframes. However, while the fixpoints of **Frith** are still easy to describe, the same does not happen with those of **BiFrm**. We leave it as an open problem to describe the categorical equivalence underlying this adjunction.

Before proceeding, we prove the following technical result:

Lemma 3.4 *Let (L, S) be a Frith frame and $a \in L$. Then, ∇_a is a positive bicomplemented element of $(C_S L, \nabla L, \Delta S)$ if and only if a is a strongly exact meet of elements of S .*

Proof Since $C_S L$ is a subframe of CL , we have that ∇_a is bicomplemented if and only if $\Delta_a \in \Delta S$, that is, if and only if there is some $P \subseteq S$ such that $\Delta_a = \bigvee_{s \in P} \Delta_s$. But this is the same as saying that $\bigwedge P$ is a strongly exact meet and $a = \bigwedge P$. \square

We are now able to prove that $\mathbf{Sk}_{\text{Frith}}$ is indeed the left adjoint of \mathbf{B}_+ .

Lemma 3.5 *The functor $\mathbf{Sk}_{\text{Frith}}$ is left adjoint to $\mathbf{B}_+ : \mathbf{BiFrm} \rightarrow \mathbf{Frith}$.*

Proof It follows from Lemma 3.4 that $\mathbf{B}_+ \circ \mathbf{Sk}_{\text{Frith}}(L, S) = (\nabla L, \{\nabla_a \mid a \in [S]_{se}\})$ and thus, the isomorphism $\nabla : L \rightarrow \nabla L$ induces an embedding of Frith frames

$$\eta_{(L,S)} : (L, S) \rightarrow \mathbf{B}_+ \circ \mathbf{Sk}_{\text{Frith}}(L, S).$$

To conclude that $\mathbf{Sk}_{\text{Frith}} \dashv \mathbf{B}_+$, it suffices to show that $\eta_{(L,S)}$ is universal from (L, S) to \mathbf{B}_+ , that is, that for every biframe \mathcal{K} and every morphism $h : (L, S) \rightarrow \mathbf{B}_+(\mathcal{K})$, there exists a unique $h' : \mathbf{Sk}_{\text{Frith}}(L, S) \rightarrow \mathcal{K}$ such that $\mathbf{B}_+(h') \circ \eta_{(L,S)} = h$. But the underlying frame homomorphism of such an h' has to be an extension $h' : C_S L \rightarrow K$ of h . Since $h[S]$ consists of complemented elements of K , by Proposition 2.1, there exists exactly one such morphism, which is easily seen to define a biframe homomorphism $h' : \mathbf{Sk}_{\text{Frith}}(L, S) \rightarrow \mathcal{K}$. \square

Corollary 3.6 *The fixpoints of **Frith** for the adjunction $\mathbf{Sk}_{\text{Frith}} \dashv \mathbf{B}_+$ are the strongly exact Frith frames.*

Proof It follows from the proof of Lemma 3.5 that the unit of the adjunction $\mathbf{Sk}_{\text{Frith}} \dashv \mathbf{B}_+$ is

$$\eta_{(L,S)} : (L, S) \rightarrow (\nabla L, \{\nabla_a \mid a \in [S]_{se}\}), \quad a \mapsto \nabla_a.$$

Clearly, this is an isomorphism if and only if (L, S) is strongly exact. \square

Let $\mathcal{L} = (L, L_+, L_-)$ be a biframe and $\mathbf{B}_+(\mathcal{L}) = (M, T)$. Since the elements of T are complemented in L , by Proposition 2.1, the frame embedding $M \hookrightarrow L$ may be uniquely extended to a frame homomorphism $C_T M \rightarrow L$. It is easily seen that this map induces a biframe homomorphism $\varepsilon_{\mathcal{L}} : (C_T M, \nabla M, \Delta T) \rightarrow (L, L_+, L_-)$. We leave it for the reader to verify that $\varepsilon_{\mathcal{L}}$ is the component at \mathcal{L} of the counit of the adjunction $\mathbf{Sk}_{\text{Frith}} \dashv \mathbf{B}_+$. In particular, if (L, S) is a Frith frame then, since $\{\Delta_s \mid s \in S\}$ and $\{\Delta_a \mid a \in [S]_{se}\}$ generate the same subframe of CL , by Lemma 3.4, we have that $\mathbf{Sk}_{\text{Frith}} \circ \mathbf{B}_+ \circ \mathbf{Sk}_{\text{Frith}}(L, S) = \mathbf{Sk}_{\text{Frith}}(L, S)$ and $\varepsilon_{\mathbf{Sk}_{\text{Frith}}(L,S)}$ is the identity map. Therefore, the adjunction $\mathbf{Sk}_{\text{Frith}} \dashv \mathbf{B}_+$ is idempotent and, as such, it induces an equivalence between the images of the two involved functors.

While it is clear that every biframe of the form $\mathbf{Sk}_{\text{Frith}}(L, S)$ is zero-dimensional, it is not the case that every zero-dimensional biframe is of that form.

Example 3.7 ⁵ Let X be a topological space such that the congruence frame of its frame of opens is not spatial (see [16, Theorem 3.4] for a characterization of the frames whose congruence frame is not spatial). We let $\mathcal{L} = (L, L_+, L_-)$ be the *Skula biframe* of X , that is: L_+ is the frame of opens of X , L_- is the subframe of $\mathcal{P}(X)$ generated by the complements of the elements of L_+ , and L is the subframe of $\mathcal{P}(X)$ generated by $L_+ \cup L_-$. To show that \mathcal{L} is not a fixpoint of the adjunction $\mathbf{Sk}_{\text{Frith}} \dashv \mathbf{B}_+$, we first recall that the underlying frame homomorphism of the counit of $\mathbf{Sk}_{\text{Frith}} \dashv \mathbf{B}_+$ at a biframe $\mathcal{L} = (L, L_+, L_-)$ is the unique frame extension $\varepsilon_{\mathcal{L}} : \mathcal{C}_T M \rightarrow L$ of the embedding $M \hookrightarrow L$, where $(M, T) = \mathbf{B}_+(\mathcal{L})$. Since, in this case, we have $\mathbf{B}_+(\mathcal{L}) = (L_+, L_+)$, \mathcal{L} is a fixpoint if and only if the unique frame homomorphism $\mathcal{C}L_+ \rightarrow L$ extending $L_+ \hookrightarrow L$ is an isomorphism. But that is not the case because L is spatial and $\mathcal{C}L_+$ is not.

A concrete example is given by taking for X the real line \mathbb{R} equipped with the Euclidean topology $\Omega(\mathbb{R})$. Since the Booleanization of $\Omega(\mathbb{R})$ is a pointless nontrivial sublocale, by the characterization of [16], its congruence frame is not spatial.

Also, the fixpoints of $\mathbf{Sk}_{\text{Frith}} \dashv \mathbf{B}_+$ need not be compact as $\mathbf{Sk}_{\text{Frith}}(L, S)$ is not compact if neither is L . We may however show that every compact and zero-dimensional biframe is a fixpoint.

Proposition 3.8 *Let $\mathcal{L} = (L, L_+, L_-)$ be a biframe and $\varepsilon_{\mathcal{L}} : (\mathcal{C}_T M, \nabla M, \Delta T) \rightarrow (L, L_+, L_-)$ be the component at \mathcal{L} of the counit of the adjunction $\mathbf{Sk}_{\text{Frith}} \dashv \mathbf{B}_+$, where $(M, T) = \mathbf{B}_+(\mathcal{L})$. Then,*

- (a) $\varepsilon_{\mathcal{L}}$ is dense,
- (b) if \mathcal{L} is zero-dimensional, then $\varepsilon_{\mathcal{L}}[\nabla M] = L_+$ and $\varepsilon_{\mathcal{L}}[\Delta T] = L_-$.

In particular, if \mathcal{L} is compact and zero-dimensional, then $\varepsilon_{\mathcal{L}}$ is an isomorphism and thus, \mathcal{L} is a fixpoint of the adjunction $\mathbf{Sk}_{\text{Frith}} \dashv \mathbf{B}_+$.

Proof Let $a \in M$ and $t \in T$ be such that $\varepsilon_{\mathcal{L}}(\nabla_a \wedge \Delta_t) = 0$. By definition of $\varepsilon_{\mathcal{L}}$, this is the same as having that the equality $a \wedge t^* = 0$ holds in L . Since t is complemented in L , this is equivalent to $a \leq t$ which, in turn, implies $\nabla_a \wedge \Delta_t = 0$. This proves that $\varepsilon_{\mathcal{L}}$ is dense.

Now, again by definition of $\varepsilon_{\mathcal{L}}$, we have that $\varepsilon_{\mathcal{L}}[\nabla M] \supseteq T$ and $\varepsilon_{\mathcal{L}}[\Delta T] \supseteq T^*$. Since T is the lattice of bicomplemented elements of L_+ , if \mathcal{L} is zero-dimensional, this implies that $\varepsilon_{\mathcal{L}}[\nabla M] = L_+$ and $\varepsilon_{\mathcal{L}}[\Delta T] = L_-$. Thus, (b) holds.

Finally, recall that $\varepsilon_{\mathcal{L}}$ is an isomorphism of biframes provided its underlying frame homomorphism is injective and satisfies $\varepsilon_{\mathcal{L}}[\nabla M] = L_+$ and $\varepsilon_{\mathcal{L}}[\Delta T] = L_-$. Thus, it suffices to show that if \mathcal{L} is compact and zero-dimensional then $\varepsilon_{\mathcal{L}}$ is injective. But it is well-known that dense frame homomorphisms with zero-dimensional domain (which is the case of $\mathcal{C}_T M$) and compact codomain are injective (see e.g. [18, Chapter VII, Proposition 2.2.2]).⁶ \square

The following is as close as we will get to a pointfree version of the result stated in Corollary 3.3.


Corollary 3.9 *Strongly exact Frith frames are a full coreflective subcategory of the category of zero-dimensional biframes.*

⁵ This example was borrowed from [7, Example 5.13] The interested reader may show that the fixpoints of the adjunction $\mathbf{Sk}_{\text{Frith}} \dashv \mathbf{B}_+$ are precisely the underlying biframes of the quasi-uniform frames representable by a Frith frame in the sense of [7, Proposition 5.6].

⁶ The result cited is stated for a *regular* domain, but every zero-dimensional frame is *regular*.

We finish this section by relating the point-set and pointfree versions of the functors we have considered.

Proposition 3.10 *The following squares commute up to natural isomorphism.*




Proof Commutativity of the left-hand side diagram follows easily from computing the functors $\mathbf{B}_+ \circ \Omega_b$ and $\Omega \circ \mathbf{Clop}_+$.

To show that the right-hand side diagram commutes up to natural isomorphism, we define a natural isomorphism $\beta : \mathbf{Sk}_{\mathbf{Pervin}} \circ \mathbf{pt} \Rightarrow \mathbf{pt}_b \circ \mathbf{Sk}_{\mathbf{Frith}}$ as follows. For a Frith frame (L, S) , we define

$$\beta_{(L,S)} : \mathbf{Sk}_{\mathbf{Pervin}} \circ \mathbf{pt}(L, S) \rightarrow \mathbf{pt}_b \circ \mathbf{Sk}_{\mathbf{Frith}}(L, S), \quad p \mapsto \tilde{p}$$

where \tilde{p} is the unique morphism making the following diagram commute (cf. Proposition 2.1).




By uniqueness of each \tilde{p} , this correspondence establishes a bijection between the points of L and the points of $C_S L$. Let us show that this is a homeomorphism with respect to the first topology of $\mathbf{Sk}_{\mathbf{Pervin}} \circ \mathbf{pt}(L, S)$. We first note that the positive open subsets of $\mathbf{Sk}_{\mathbf{Pervin}} \circ \mathbf{pt}(L, S)$ are the subsets of the form \widehat{a} , while those of $\mathbf{pt}_b \circ \mathbf{Sk}_{\mathbf{Frith}}(L, S)$ are the subsets of the form $\widehat{\nabla a}$, where $a \in L$. Now, given $a \in L$ and $p \in \mathbf{pt}(L)$, using commutativity of the triangle above, we have

$$p \in \beta_{(L,S)}^{-1}(\widehat{\nabla a}) \iff \tilde{p}(\nabla a) = 1 \iff p(a) = 1 \iff p \in \widehat{a}.$$

Since $\beta_{(L,S)}$ is a bijection, this implies that $\beta_{(L,S)}$ is both continuous and open with respect to the first topologies, thus a homeomorphism. Showing that $\beta_{(L,S)}$ is also a homeomorphism with respect to the second topology is analogous, and we leave it for the reader.

Now, β is a natural transformation provided the following square commutes for every morphism $h : (M, T) \rightarrow (L, S)$ of Frith frames.



That is indeed the case because, for every $p \in \text{pt}(L)$ and $x \in M$, we have the following equalities:

$$\tilde{p} \circ \mathbf{Sk}_{\text{Frith}}(h)(\nabla_x) = \tilde{p}(\nabla_{h(x)}) = p \circ h(x) = \widetilde{p \circ h}(\nabla_x).$$

□

Proposition 3.10 makes it natural to ask whether the corresponding diagrams for the spectrum and open-set functors also commute. The answer is negative, as shown by the next example.

Example 3.11 For the first diagram, consider the biframe $\mathcal{L} = (\mathbf{3}, \mathbf{3}, \mathbf{2})$, where $\mathbf{3}$ denotes the 3-element chain. Then, $\text{pt} \circ \mathbf{B}_+(\mathcal{L})$ is a space with one point, while $\mathbf{Clop}_+ \circ \text{pt}_b(\mathcal{L})$ has two, so these cannot be isomorphic. For the second diagram, we let X be a topological space as in Example 3.7 and observe that $\Omega_b \circ \mathbf{Sk}_{\text{Pervin}}(X, \Omega(X))$ is the *Skula biframe* \mathcal{L} of X . As already argued, \mathcal{L} is not in the image of $\mathbf{Sk}_{\text{Frith}}$, thus, $\Omega_b \circ \mathbf{Sk}_{\text{Pervin}}(X, \Omega(X))$ is not isomorphic to $\mathbf{Sk}_{\text{Frith}} \circ \Omega(X, \Omega(X))$.

3.2 The Monotopological Case

In this section, we will investigate the monotopological version of the results of Sect. 3.1. Under the identifications $\mathbf{Top} \hookrightarrow \mathbf{BiTop}$ and $\mathbf{Frm} \hookrightarrow \mathbf{BiFrm}$, these will follow as a consequence of the latter.

Let us consider the restrictions $\mathbf{Clop} : \mathbf{Top} \rightarrow \mathbf{Pervin}$ and $\mathbf{B} : \mathbf{Frm} \rightarrow \mathbf{Frith}$ of the functors \mathbf{Clop}_+ and \mathbf{B}_+ defined in Sect. 3.1. Explicitly, \mathbf{Clop} maps a topological space (X, τ) to the Pervin space $\mathbf{Clop}_+(X, \tau, \tau) = (X, \mathbf{Clop}(X, \tau))$, where $\mathbf{Clop}(X, \tau)$ denotes the Boolean algebra of clopen subsets of X for the topology τ , and a morphism to itself. On the other hand, given a frame L , $\mathbf{B}(L)$ is the pair $(\langle \mathbf{B}(L) \rangle_{\mathbf{Frm}}, \mathbf{B}(L))$, where $\langle \mathbf{B}(L) \rangle_{\mathbf{Frm}}$ denotes the subframe of L generated by the lattice of complemented elements $\mathbf{B}(L)$ of L , and a frame homomorphism $h : L \rightarrow K$ is sent to the morphism of Frith frames $\mathbf{B}h : \mathbf{B}(L) \rightarrow \mathbf{B}(K)$ induced by the suitable restriction and co-restriction of h . Then, the adjunctions $\mathbf{Clop}_+ : \mathbf{BiTop} \rightleftarrows \mathbf{Pervin} : \mathbf{Sk}_{\text{Pervin}}$ and $\mathbf{Sk}_{\text{Frith}} : \mathbf{Frith} \rightleftarrows \mathbf{BiFrm} : \mathbf{B}_+$ studied in Sect. 3.1 restrict, respectively, to adjunctions $\mathbf{Clop} : \mathbf{Top} \rightleftarrows \mathbf{Pervin}' : \mathbf{U}'_{\text{Pervin}}$ and $\mathbf{U}'_{\text{Frith}} : \mathbf{Frith}' \rightleftarrows \mathbf{Frm} : \mathbf{B}$, where

- \mathbf{Pervin}' denotes the full subcategory of \mathbf{Pervin} determined by the Pervin spaces (X, \mathcal{S}) such that $\mathbf{Sk}_{\text{Pervin}}(X, \mathcal{S})$ belongs to the image of $\mathbf{Top} \hookrightarrow \mathbf{BiTop}$,
- \mathbf{Frith}' denotes the full subcategory of \mathbf{Frith} determined by the Frith frames (L, \mathcal{S}) such that $\mathbf{Sk}_{\text{Frith}}(L, \mathcal{S})$ belongs to the image of $\mathbf{Frm} \hookrightarrow \mathbf{BiFrm}$,
- $\mathbf{U}'_{\text{Pervin}}$ is the suitable restriction and co-restriction of $\mathbf{Sk}_{\text{Pervin}}$, and
- $\mathbf{U}'_{\text{Frith}}$ is the suitable restriction and co-restriction of $\mathbf{Sk}_{\text{Frith}}$.

The following result, whose proof is trivial, explains our choice of notation for the functors $\mathbf{U}'_{\text{Pervin}}$ and $\mathbf{U}'_{\text{Frith}}$: these are nothing but the suitable restrictions of the functors $\mathbf{U}_{\text{Pervin}}$ and $\mathbf{U}_{\text{Frith}}$ defined in Sect. 2.6.

Lemma 3.12 *The following statements hold:*

- (a) a Pervin space (X, \mathcal{S}) belongs to \mathbf{Pervin}' if and only if $\Omega_{\mathcal{S}}(X) = \Omega_{\mathcal{S}^c}(X)$,
- (b) a Frith frame (L, \mathcal{S}) belongs to \mathbf{Frith}' if and only if $\nabla L = \Delta \mathcal{S}$.

In particular, given $(X, \mathcal{S}) \in \mathbf{Pervin}'$ and $(L, \mathcal{S}) \in \mathbf{Frith}'$, the following equalities hold:

$$\mathbf{U}'_{\text{Pervin}}(X, \mathcal{S}) = (X, \Omega_{\mathcal{S}}(X)) \quad \text{and} \quad \mathbf{U}'_{\text{Frith}}(L, \mathcal{S}) = L.$$

We will now characterize the categorical equivalences induced by the adjunctions $\mathbf{Clop} \dashv \mathbf{U}'_{\text{Pervin}}$ and $\mathbf{U}'_{\text{Frith}} \dashv \mathbf{B}$. Recall that we have seen in Sect. 3.1 that both $\mathbf{Clop}_+ \dashv \mathbf{Sk}_{\text{Pervin}}$ and $\mathbf{Sk}_{\text{Frith}} \dashv \mathbf{B}_+$ are idempotent and, therefore, so are their restrictions.

Proposition 3.13 *The categories of zero-dimensional topological spaces and that of strongly exact symmetric Pervin spaces are equivalent.*

Proof Since $\mathbf{Clop} \dashv \mathbf{U}'_{\text{Pervin}}$ is a restriction of $\mathbf{Clop}_+ \dashv \mathbf{Sk}_{\text{Pervin}}$, by Proposition 3.2, it induces an equivalence between the categories of topological spaces that are zero-dimensional when seen as bitopological spaces, and the category determined by the Pervin spaces $(X, S) \in \mathbf{Pervin}'$ that are strongly exact. The former are easily seen to be the zero-dimensional topological spaces. We argue that $(X, S) \in \mathbf{Pervin}'$ is strongly exact if and only if it is a strongly exact symmetric Pervin space. Clearly, \mathbf{Pervin}' contains all symmetric Pervin spaces, thus the backwards implication is trivial. Conversely, if $(X, S) \in \mathbf{Pervin}'$ is strongly exact then, being a fixpoint of the idempotent adjunction $\mathbf{Clop} \dashv \mathbf{U}'_{\text{Pervin}}$, it belongs to the image of \mathbf{Clop} . Hence, it is symmetric, as required. \square

Proposition 3.14 *The categories of zero-dimensional frames and that of strongly exact symmetric Frith frames are equivalent.*

Proof Since the adjunction $\mathbf{U}'_{\text{Frith}} \dashv \mathbf{B}$ is idempotent, its fixpoints in \mathbf{Frm} are the frames of the form $\mathbf{U}'_{\text{Frith}}(L, S) = L$, for $(L, S) \in \mathbf{Frith}'$. Noticing that the inclusion $\nabla L \supseteq \Delta S$ holds if and only if S consists of complemented elements, these are easily seen to be the zero-dimensional ones. On the other hand, an argument similar to that used in the proof of Proposition 3.13 shows that the fixpoints of $\mathbf{U}'_{\text{Frith}} \dashv \mathbf{B}$ in \mathbf{Frith}' are strongly exact symmetric Frith frames. \square

4 Complete Pervin Spaces and Complete Frith Frames

In this section we will show that the dual adjunction $\mathbf{\Omega} : \mathbf{Pervin} \rightleftharpoons \mathbf{Frith} : \mathbf{pt}$ induces a duality between T_0 complete Pervin spaces on the one hand and complete Frith frames on the other. We use the following definition from [11, 19].

Definition 4.1 ([11, 19]) Let (X, S) be a Pervin space. A filter $F \subseteq \mathcal{P}(X)$ is a *Cauchy filter* if it is proper and, for every $S \in \mathcal{S}$, either S or its complement is in F . We say that a Cauchy filter F *converges* to the point $x \in X$ if every open neighborhood $U \in \Omega_{\overline{S}}(X)$ of x belongs to F . Finally, a Pervin space (X, S) is said to be *Cauchy complete* if every Cauchy filter converges, and a *Cauchy completion* of (X, S) is a dense extremal monomorphism $c : (X, S) \hookrightarrow (Y, T)$ into a Cauchy complete Pervin space (Y, T) .⁷

In the following, we refer to the symmetrization of a Pervin space, defined in Sect. 2.5. The following is an easy observation that we state for later reference.

Lemma 4.2 *Let (X, S) be a Pervin space, and $F \subseteq \mathcal{P}(X)$ be a Cauchy filter. Then, F converges to x if and only if x belongs to $\bigcap(F \cap \overline{S})$.*

Note that a filter is Cauchy with respect to (X, S) if and only if it is Cauchy with respect to (X, \overline{S}) . Therefore, a Pervin space is Cauchy complete if and only if so is its symmetrization.

⁷ In [11] *Cauchy complete* and *Cauchy completion* are simply named *complete* and *completion*, respectively.

As observed in [11, 19], one may show that this notion of Cauchy complete Pervin space correctly captures the notion of a complete quasi-uniform space. It is known that complete quasi-uniform spaces may be equivalently characterized via dense extremal monomorphisms. In the case of Pervin spaces, the suitable definitions are the following.

Definition 4.3 A symmetric Pervin space (X, \mathcal{B}) is *complete* if every dense extremal monomorphism $(X, \mathcal{B}) \hookrightarrow (Y, \mathcal{C})$, with (Y, \mathcal{C}) a T_0 symmetric Pervin space is an isomorphism. More generally, we say that a Pervin space (X, \mathcal{S}) is *complete* if so is its symmetrization.

Our next goal is to show that Definitions 4.1 and 4.3 are equivalent. Before we move on, we need to prove a couple of technical lemmas.

Lemma 4.4 *Let $m : (X, \mathcal{S}) \hookrightarrow (Y, \mathcal{T})$ be a dense extremal monomorphism. If $F \subseteq \mathcal{P}(Y)$ is a Cauchy filter, then so is $m^{-1}(F)$.*

Proof Let $F \subseteq \mathcal{P}(Y)$ be a Cauchy filter. Since m is dense and F is proper, $m^{-1}(F)$ is, by Lemma 2.9, a proper filter too. Now, given $S \in \mathcal{S}$, since m is an extremal monomorphism, we have $S = m^{-1}(T)$ for some $T \in \mathcal{T}$. Since F is Cauchy, it contains either T or T^c and thus, $m^{-1}(F)$ contains either $S = m^{-1}(T)$ or $S^c = m^{-1}(T^c)$. This shows that $m^{-1}(F)$ is Cauchy as well. □

Recall the neighborhood map $\mathcal{N}_{(X, \mathcal{S})} : (X, \mathcal{S}) \rightarrow (\text{pf}(\mathcal{S}), \tilde{\mathcal{S}})$ from (5), that is, the unit of the adjunction $\mathbf{L}_{\text{Pervin}} \dashv \mathbf{pf}$.

Lemma 4.5 *For a T_0 Pervin space (X, \mathcal{S}) , the map $\mathcal{N}_{(X, \mathcal{S})}$ is an extremal monomorphism of Pervin spaces whose symmetrization is dense.*

Proof If (X, \mathcal{S}) is T_0 , then different points have different neighborhood filters in \mathcal{S} , and so $\mathcal{N}_{(X, \mathcal{S})}$ is injective. Since, for every $S \in \mathcal{S}$, we have $\mathcal{N}_{(X, \mathcal{S})}^{-1}(\tilde{S}) = S$, the map $\mathcal{N}_{(X, \mathcal{S})}$ is an extremal monomorphism. To show that $\mathbf{Sym}_{\text{Perv}}(\mathcal{N}_{(X, \mathcal{S})})$ is dense, suppose that $\mathcal{N}_{(X, \mathcal{S})}^{-1}(\tilde{S}_1 \cap \tilde{S}_2^c) = S_1 \cap S_2^c = \emptyset$, that is, $S_1 \subseteq S_2$. Then, there is no prime filter containing S_1 and omitting S_2 , which means that $\tilde{S}_1 \cap \tilde{S}_2^c$ must be empty. □

We remark that, for every T_0 Pervin space (X, \mathcal{S}) , the map $\mathcal{N}_{(X, \mathcal{S})} : (X, \mathcal{S}) \hookrightarrow (\text{pf}(\mathcal{S}), \tilde{\mathcal{S}})$ is the completion of (X, \mathcal{S}) (cf. [11, 19]).

We may now prove the following characterization of T_0 complete Pervin spaces.

Theorem 4.6 *Let (X, \mathcal{S}) be a T_0 Pervin space. Then, the following are equivalent:*

- (a) (X, \mathcal{S}) is Cauchy complete;
- (b) (X, \mathcal{S}) is complete;
- (c) Every extremal monomorphism $(X, \mathcal{S}) \hookrightarrow (Y, \mathcal{T})$ into a T_0 Pervin space whose symmetrization is dense is an isomorphism;
- (d) (X, \mathcal{S}) is isomorphic to $\mathbf{pf}(\mathcal{S})$;
- (e) (X, \mathcal{S}) is isomorphic to $\mathbf{pt}(\text{Idl}(\mathcal{S}), \mathcal{S})$;
- (f) (X, \mathcal{S}) is isomorphic to a Pervin space of the form $\mathbf{pt}(\text{Idl}(D), D)$, for some lattice D ;
- (g) (X, \mathcal{S}) is isomorphic to a Pervin space of the form $\mathbf{pf}(D)$, for some lattice D .

Proof Noting that extremal monomorphisms are preserved under symmetrization, the equivalence between (b) and (c) follows. That (c) implies (d) is a consequence of Lemma 4.5. The

equivalences between (d) and (e) and between (f) and (g) follow from Lemma 2.11, while that (e) implies (f) is trivial. It remains to show that (a) implies (b) and that (g) implies (a).

Suppose that (a) holds, and let $m : (X, \overline{S}) \hookrightarrow (Y, C)$ be a dense extremal monomorphism into a symmetric T_0 Pervin space (Y, C) . We need to show that m is an isomorphism, that is, that m is surjective. Given $y \in Y$, consider the filter $F_y := \uparrow\{C \in C \mid y \in C\}$. Since C is a Boolean algebra, F_y is a Cauchy filter. By Lemma 4.4, the filter $m^{-1}(F_y)$ is Cauchy as well. Since (X, S) is Cauchy complete, $m^{-1}(F_y)$ converges to some point $x \in X$. We claim that $y = m(x)$. Since (Y, C) is T_0 and C is a Boolean algebra, we have that $y = m(x)$ provided $y \in C$ implies $m(x) \in C$, for every $C \in C$. We let $C \in C$ be such that $y \in C$. Then, $m^{-1}(C)$ belongs to $m^{-1}(F_y) \cap \overline{S}$ and, since $m^{-1}(F_y)$ converges to x , by Lemma 4.2, it follows that $x \in m^{-1}(C)$, that is, $m(x) \in C$, as required.

Finally, let us assume that we have a Pervin space of the form $\mathbf{pf}(D) = (\mathbf{pf}(D), \widetilde{D})$, with D a distributive lattice. Suppose that $F \subseteq \mathcal{P}(\mathbf{pf}(D))$ is a Cauchy filter and set $P := \{a \in D \mid \widetilde{a} \in F\}$. Clearly, P is a filter of D . Let us show that P is prime. Let $a, b \in D$ be such that $a \vee b \in P$, and suppose that $a \notin P$. Equivalently, a, b are such that $\widetilde{a \vee b} = \widetilde{a} \cup \widetilde{b} \in F$ and $\widetilde{a} \notin F$. Since F is Cauchy, it follows that $(\widetilde{a})^c$ belongs to F and thus, so does $\widetilde{b} \supseteq (\widetilde{a})^c \cap \widetilde{b} = (\widetilde{a})^c \cap (\widetilde{a} \cup \widetilde{b})$. This shows that $b \in P$ as required. We now claim that F converges to P . By Lemma 4.2, it suffices to show that $P \in \widetilde{a} \cap (\widetilde{b})^c$ whenever $a, b \in D$ are such that $\widetilde{a} \cap (\widetilde{b})^c \in F$. Since F is proper, having $\widetilde{a} \cap (\widetilde{b})^c \in F$ implies that $\widetilde{a} \in F$ and $\widetilde{b} \notin F$. But by definition of P , this means that $P \in \widetilde{a} \cap (\widetilde{b})^c$, as required. \square

From now on, we will drop the use of *Cauchy complete* and call *complete Pervin space* every Pervin space satisfying the equivalent conditions of Theorem 4.6.

In turn, completeness of Frith frames is discussed in [7], where a characterization of complete Frith frames using both dense extremal epimorphisms and Cauchy maps is given. Here, it will suffice to consider the following definitions:

Definition 4.7 ([7]) We say that a symmetric Frith frame (L, B) is *complete* if every dense extremal epimorphism $(M, C) \twoheadrightarrow (L, B)$ with (M, C) symmetric is an isomorphism. More generally, a Frith frame (L, S) is *complete* provided its symmetric reflection $\mathbf{Sym}_{\mathbf{Frith}}(L, S)$ is complete. A *completion* of (L, S) is a complete Frith frame (M, T) together with a dense extremal epimorphism $(M, T) \twoheadrightarrow (L, S)$.

The fact that completeness of a Frith frame (L, S) is equivalent to completeness of the associated quasi-uniform frame is shown in [7, Proposition 7.2]. Moreover, every Frith frame (L, S) has a unique, up to isomorphism, completion, which is given by the counit (3) of the adjunction $\mathbf{Idl} \dashv \mathbf{L}_{\mathbf{Frith}}$. Also in [7], we have shown the following:

Theorem 4.8 ([7, Proposition 4.6 and 0 Theorem 7.7]) *Let (L, S) be a Frith frame. Then, the following are equivalent:*

- (a) (L, S) is complete;
- (b) (L, S) is coherent;
- (c) L is isomorphic to the ideal completion $\mathbf{Idl}(S)$ of S .

Part (c) of this characterization, together with Theorem 4.6(f), yield the following:

Corollary 4.9 *A Pervin space is T_0 and complete if and only if it is of the form $\mathbf{pt}(L, S)$, for some complete Frith frame (L, S) .*

In particular, the functor $\mathbf{pt} : \mathbf{Frith} \rightarrow \mathbf{Pervin}$ restricts and co-restricts to a functor $\mathbf{CFrith} \rightarrow \mathbf{CPervin}$. In order to show that Ω , too, restricts correctly, we will need to use the Prime Ideal Theorem.

Proposition 4.10 *The following are equivalent.*

- (a) *The Prime Ideal Theorem holds;*
- (b) *If (X, S) is a T_0 complete Pervin space, then the Frith frame $\Omega(X, S)$ is complete.*
- (c) *If (X, \mathcal{B}) is a T_0 complete symmetric Pervin space, then the Frith frame $\Omega(X, \mathcal{B})$ is complete.*

Proof We first show that (a) implies (b). Let (X, S) be a T_0 complete Pervin space. By Theorem 4.6, we may assume, without loss of generality, that $(X, S) = \mathbf{pt}(\text{Idl}(S), S)$. Since we are assuming that the Prime Ideal Theorem holds, by Theorem 2.2, $\text{Idl}(S)$ is a spatial frame, and thus so is $(\text{Idl}(S), S)$ (cf. Theorem 2.5). Therefore, we have

$$\Omega(X, S) = \Omega \circ \mathbf{pt}(\text{Idl}(S), S) \cong (\text{Idl}(S), S),$$

which, by Theorem 4.8, is a complete Frith frame.

Clearly, (b) implies (c). Finally, suppose that (c) holds, and let X be a set. By Theorem 2.2, it suffices to show that every proper filter F on X is contained in a prime filter. For the sake of readability, we set $B := \mathcal{P}(X)$. Note that showing that F is contained in some prime filter is equivalent to showing that the intersection $\bigcap_{S \in F} \tilde{S}$ is nonempty. For that, we consider the T_0 symmetric Pervin space $(\text{pf}(B), \tilde{B})$. By Theorem 4.6, this is complete, and by hypothesis, so is the Frith frame $\Omega(\text{pf}(B), \tilde{B}) = (\Omega_{\tilde{B}}(\text{pf}(B)), \tilde{B})$. In particular, the topological space $(\text{pf}(B), \Omega_{\tilde{B}}(\text{pf}(B)))$ is compact, and thus, it suffices to show that $\{\tilde{S} \mid S \in F\}$ has the finite intersection property. In turn, since F is closed under finite meets, this is the same as showing that $\tilde{S} \neq \emptyset$ for every $S \in F$. But since F is proper, every $S \in F$ is nonempty, and given $x \in S$, we have $\uparrow\{x\} \in \tilde{S}$ and \tilde{S} is nonempty as well. This finishes the proof. \square

Corollary 4.11 *The following are equivalent.*

- (a) *The Prime Ideal Theorem holds;*
- (b) *If (X, S) is a T_0 complete Pervin space, then the compact open subsets of $(X, \Omega_S(X))$ are precisely the elements of S .*

Proof By the equivalence between (a) and (b) of Proposition 4.10, it suffices to show that $\Omega(X, S)$ is a complete Frith frame if and only if the compact open subsets of $(X, \Omega_S(X))$ are precisely the elements of S . Observe that, by Theorem 4.8, $\Omega(X, S) = (\Omega_S(X), S)$ is complete if and only if it is coherent. Thus, S is the set of compact elements of the frame $\Omega_S(X)$, hence of compact open subsets of the topological space $(X, \Omega_S(X))$. \square

We have just proved the following:

Corollary 4.12 *If the Prime Ideal Theorem holds, then the adjunction $\Omega : \mathbf{Pervin} \rightleftarrows \mathbf{Frith} : \mathbf{pt}$ restricts and co-restricts to an adjunction between T_0 complete Pervin spaces and complete Frith frames.*

We may now show the main result of this section.

Theorem 4.13 *If the Prime Ideal Theorem holds, then the adjunction $\Omega : \mathbf{Pervin} \rightleftarrows \mathbf{Frith} : \mathbf{pt}$ restricts and co-restricts to a duality between the categories $\mathbf{CPervin}$ of T_0 complete Pervin spaces and the category \mathbf{CFrith} of complete Frith frames.*

Proof Recall from Theorem 2.5 that the adjunction $\Omega \dashv \mathbf{pt}$ induces a duality between sober Pervin spaces and spatial Frith frames. Thus, because of Corollary 4.12, it remains to show that T_0 complete Pervin spaces are sober and complete Frith frames are spatial. By Theorem 4.6, a

T_0 Pervin space is complete if and only if it is isomorphic to the Pervin space $\mathbf{pt}(\text{Idl}(D), D)$, for some lattice D , and thus a complete Pervin space is sober. In turn, complete Frith frames are spatial because, by Theorem 4.8, they are of the form $(\text{Idl}(D), D)$ for some lattice D and, by the Prime Ideal Theorem, these are spatial Frith frames (cf. Theorems 2.2 and 2.5). \square

5 Stone-Type Dualities

In this section we will see how several Stone-type dualities relate to the duality between T_0 complete Pervin spaces and complete Frith frames shown in the previous section (cf. Theorem 4.13). Note that the fact that every T_0 complete Pervin space defines a spectral, a Priestley, and a pairwise Stone space is already observed in [19], but no proof is provided. Here, we will give the functorial details of this assignment, and use Theorem 4.13 to interpret Stone-type dualities as a restriction and co-restriction of the dual adjunction $\Omega : \mathbf{Pervin} \rightleftarrows \mathbf{Frith} : \mathbf{pt}$ along *full* subcategory embeddings.

5.1 Stone Duality

Stone duality establishes that the categories of bounded distributive lattices and of spectral spaces are dually equivalent. We recall that a topological space is *spectral* if it is sober, and its compact open subsets are closed under finite intersections and form a basis of the topology. The category of spectral spaces together with those continuous functions such that the preimages of compact open subsets are compact will be denoted by \mathbf{Spec} . By identifying each lattice with the coherent frame given by its ideal completion, Stone duality may be seen as a restriction and co-restriction of the dual adjunction $\Omega : \mathbf{Top} \rightleftarrows \mathbf{Frm} : \mathbf{pt}$, but *not along full inclusions*, as not every continuous function is a morphism of spectral spaces, and morphisms of coherent frames are required to preserve compact elements. We will now see that this duality may also be seen as a restriction and co-restriction of the dual adjunction $\Omega : \mathbf{Pervin} \rightleftarrows \mathbf{Frith} : \mathbf{pt}$, the advantage being that spectral spaces and bounded distributive lattices form *full* subcategories of \mathbf{Pervin} and \mathbf{Frith} , respectively.

It follows straightforwardly from Theorem 4.8 (by identifying each lattice S with the Frith frame $(\text{Idl}(S), S)$) that the categories of complete Frith frames and of bounded distributive lattices are equivalent, and by definition of morphism of Frith frames, this is indeed a full subcategory of \mathbf{Frith} (see also [7, Proposition 4.6]). Given the Prime Ideal Theorem, we also have that every T_0 complete Pervin space defines a spectral space.

Lemma 5.1 *If the Prime Ideal Theorem holds, then the functor $\mathbf{U}_{\text{Pervin}} : \mathbf{Pervin} \rightarrow \mathbf{Top}$ restricts and co-restricts to a functor $\mathbf{CPervin} \rightarrow \mathbf{Spec}$.*

Proof Let (X, S) be a T_0 complete Pervin space. By Theorem 4.13, (X, S) is a fixpoint of the adjunction $\Omega \dashv \mathbf{pt}$ and thus, it is sober, that is to say that the topological space $(X, \Omega_S(X))$ is sober (recall Theorem 2.5). By Corollary 4.11, the lattice S is the set of compact open elements of $(X, \Omega_S(X))$. Therefore, $(X, \Omega_S(X))$ is spectral. Finally, note that this also implies that if $f : (X, S) \rightarrow (Y, T)$ is a morphism of T_0 complete Pervin spaces, then it induces a morphism $f : (X, \Omega_S(X)) \rightarrow (Y, \Omega_T(Y))$ of spectral spaces. \square

It remains then to show that the categories of T_0 complete Pervin spaces and of spectral spaces are, in fact, isomorphic. Consider the functor $\mathbf{KO} : \mathbf{Spec} \rightarrow \mathbf{CPervin}$ which sends a spectral space (X, τ) to the Pervin space X equipped with the lattice of compact open subsets of X . It is easily seen that \mathbf{KO} is well-defined. We may further prove the following:

Proposition 5.2 *If the Prime Ideal Theorem holds, then the functors $\mathbf{U}_{\text{Pervin}}$ and \mathbf{KO} establish an isomorphism of categories between \mathbf{Spec} and $\mathbf{CPervin}$.*

Proof Let (X, \mathcal{S}) be a T_0 complete Pervin space. By Corollary 4.11, $\mathbf{KO} \circ \mathbf{U}_{\text{Pervin}}$ is the identity functor on $\mathbf{CPervin}$. For a spectral space (X, τ) , we have that $\mathbf{U}_{\text{Pervin}} \circ \mathbf{KO}(X, \tau)$ is the set X equipped with the topology generated by the lattice of compact open subsets of (X, τ) . But by definition of spectral space, this is (X, τ) itself. \square

Corollary 5.3 *If the Prime Ideal Theorem holds, then Stone duality for bounded distributive lattices may be seen as a restriction along full subcategory embeddings of the dual adjunction $\Omega : \mathbf{Pervin} \rightleftarrows \mathbf{Frith} : \mathbf{pt}$.*

5.2 Priestley Duality

Another duality for bounded distributive lattices, due to Priestley, uses the so-called *Priestley spaces* in place of spectral spaces. A *Priestley space* is a compact topological space equipped with a partial order relation on its points satisfying the *Priestley separation axiom*, which states that for points $x \not\leq y$ there is a clopen upper set (“upset” hereon) containing x and omitting y . A morphism of Priestley spaces is a continuous map which is monotone with respect to the order. We denote by \mathbf{Priest} the category of Priestley spaces and corresponding morphisms. It has been shown in [8] that the category of Priestley spaces and that of spectral spaces are isomorphic. We have already seen that T_0 complete Pervin spaces form a category equivalent to spectral spaces, and it is well-known that the latter form a category equivalent to \mathbf{Priest} . It is the goal of this section to explicitly exhibit the correspondence between T_0 complete Pervin spaces and Priestley spaces, thereby providing yet another way of understanding Priestley duality.

As understood in [19], every Pervin space (X, \mathcal{S}) comes naturally equipped with a preorder given by

$$x \leq_{\mathcal{S}} y \text{ if and only if } x \in S \text{ implies } y \in S \text{ for all } S \in \mathcal{S},$$

which is a partial order exactly when (X, \mathcal{S}) is T_0 . In the case where (X, \mathcal{S}) is T_0 and complete, this is the underlying partial order of the corresponding Priestley space, and its topology is the patch topology of the bitopological space $\mathbf{Sk}_{\text{Pervin}}(X, \mathcal{S})$.

Lemma 5.4 *If the Prime Ideal Theorem holds, then there is a well-defined functor $\mathbf{P} : \mathbf{CPervin} \rightarrow \mathbf{Priest}$ defined by $\mathbf{P}(X, \mathcal{S}) = (X, \Omega_{\overline{\mathcal{S}}}(X), \leq_{\mathcal{S}})$ on objects, and mapping each morphism to the morphism defined by its underlying set function.*

Proof If (X, \mathcal{S}) is T_0 complete, then so is its symmetrization $(X, \overline{\mathcal{S}})$ and, by Corollary 4.11, $X \in \overline{\mathcal{S}}$ is a compact element of $\Omega_{\overline{\mathcal{S}}}(X)$. Thus, $(X, \Omega_{\overline{\mathcal{S}}}(X))$ is compact. Since elements of \mathcal{S} are clopen upsets of $(X, \Omega_{\overline{\mathcal{S}}}(X), \leq_{\mathcal{S}})$, the relation $\leq_{\mathcal{S}}$ satisfies the Priestley separation axiom. Therefore, $\mathbf{P}(X, \mathcal{S})$ is a Priestley space. It is not hard to see that \mathbf{P} is well-defined on morphisms, too. \square

In the other direction, we have the following:

Lemma 5.5 *There is a well-defined functor $\mathbf{CUp} : \mathbf{Priest} \rightarrow \mathbf{CPervin}$ that assigns to each Priestley space (X, τ, \leq) the Pervin space X equipped with the lattice of clopen upsets of X , and keeps morphisms unchanged.*

Proof It is easy to verify that **CU_p** is well-defined on morphisms. Let us argue that **CU_p** is well-defined on objects. Fix a Priestley space (X, τ, \leq) . By the Priestley separation axiom, we have that $(X, \mathcal{S}) := \mathbf{CU}_p(X, \tau, \leq)$ is T_0 . To show that (X, \mathcal{S}) is complete, we show that every Cauchy filter converges. Indeed, if $F \subseteq \mathcal{P}(X)$ is a Cauchy filter then, since it is proper, it has the finite intersection property. Thus, $F \cap \overline{\mathcal{S}}$ is a family of closed subsets of X with the finite intersection property. Since Priestley spaces are compact, it follows that $\bigcap(F \cap \overline{\mathcal{S}})$ is nonempty and, by Lemma 4.2, F converges. \square

We leave it for the reader to verify that the functors **P** and **CU_p** are mutually inverse. The reader may also check that, by composing the functors

$$\mathbf{Priest} \xrightarrow{\mathbf{CU}_p} \mathbf{CPervin} \xrightarrow{U_{\text{Pervin}}} \mathbf{Spec} \quad \text{and} \quad \mathbf{Spec} \xrightarrow{\mathbf{P}} \mathbf{CPervin} \xrightarrow{\mathbf{KO}} \mathbf{Priest}$$

one obtains the well-known isomorphism between the categories of spectral and of Priestley spaces.

Proposition 5.6 *If the Prime Ideal Theorem holds, then the functors **P** and **CU_p** establish an isomorphism of categories between **Priest** and **CPervin**.*

Corollary 5.7 *If the Prime Ideal Theorem holds, then Priestley duality may be seen as a restriction along full subcategory embeddings of the dual adjunction $\mathbf{\Omega} : \mathbf{Pervin} \rightleftarrows \mathbf{Frith} : \mathbf{pt}$.*

5.3 Bitopological Duality

It has long been known [3] that the dual adjunction between bitopological spaces and biframes restricts and co-restricts to a duality between T_0 , compact and zero-dimensional bitopological spaces (denoted **BiTop_{KZ}**) and compact and zero-dimensional biframes (denoted **BiFrm_{KZ}**). A few years later, Priestley duality was considered from a bitopological point of view [17], with Priestley spaces being identified with T_0 , compact and zero-dimensional bitopological spaces, and lattices being identified with compact and zero-dimensional biframes. The point-set half of this correspondence was then rediscovered in [6], where T_0 compact zero-dimensional bitopological spaces were named *pairwise Stone spaces*.⁸ It is then clear that **CPervin** and **CFrith** are equivalent to **BiTop_{KZ}** and **BiFrm_{KZ}**, respectively. In this section, we will make these equivalences explicit, using the adjunctions derived in Sect. 3.1.

Let us start with the equivalence between T_0 complete Pervin spaces and T_0 , compact, and zero-dimensional bitopological spaces. Recall that we have an idempotent adjunction $\mathbf{Clo}_+ \dashv \mathbf{Sk}_{\text{Pervin}}$ whose fixpoints are, respectively, the zero-dimensional bitopological spaces and the strongly exact Pervin spaces. To conclude that this adjunction further restricts to an equivalence between **BiTop_{KZ}** and **CPervin**, it suffices to show that T_0 complete Pervin spaces are strongly exact (cf. Corollary 5.9) and that the equality $\mathbf{Sk}_{\text{Pervin}}[\mathbf{CPervin}] = \mathbf{BiTop}_{KZ}$ holds (cf. Lemmas 5.8 and 5.10). For that, we will need to assume the Prime Ideal Theorem.

Lemma 5.8 *The following are equivalent.*

- (a) *The Prime Ideal Theorem holds.*
- (b) *For a T_0 complete Pervin space (X, \mathcal{S}) , the bispaces $\mathbf{Sk}_{\text{Pervin}}(X, \mathcal{S})$ is compact.*

⁸ Note that pairwise Stone spaces are the same as T_0 , compact and zero-dimensional bitopological spaces only under the assumption of the Prime Ideal Theorem.

(c) For a T_0 symmetric complete Pervin space (X, \mathcal{B}) , the bispace $\mathbf{Sk}_{\text{Pervin}}(X, \mathcal{B})$ is compact.

Proof We first argue that (b) and (c) are equivalent. Clearly, (b) implies (c). For the converse, we only need to remind the reader that a Pervin space (X, \mathcal{S}) is complete if and only if so is its symmetrization $(X, \overline{\mathcal{S}})$ and observe that, by definition of compact bispace, $\mathbf{Sk}_{\text{Pervin}}(X, \mathcal{S})$ is compact if and only if so is $\mathbf{Sk}_{\text{Pervin}}(X, \overline{\mathcal{S}})$.

Now, taking the equivalence between statements (a) and (c) of Proposition 4.10 into account, it suffices to show that, for every T_0 symmetric Pervin space (X, \mathcal{B}) , the bispace $\mathbf{Sk}_{\text{Pervin}}(X, \mathcal{B})$ is compact if and only if the Frith frame $\mathbf{\Omega}(X, \mathcal{B})$ is coherent (recall that, by Theorem 4.8, a Frith frame is coherent if and only if it is complete).

If $\mathbf{Sk}_{\text{Pervin}}(X, \mathcal{B})$ is compact, that is, if the topological space $(X, \Omega_{\mathcal{B}}(X))$ is compact, then every element of \mathcal{B} is compact in the frame $\Omega_{\mathcal{B}}(X)$. Indeed, if $B = \bigcup \mathcal{U}$ for some family $\mathcal{U} \subseteq \Omega_{\mathcal{B}}(X)$, then $\mathcal{U} \cup \{B^c\}$ is an open cover of $(X, \Omega_{\mathcal{B}}(X))$ and, if the latter is compact, then $\mathcal{U} \cup \{B^c\}$ has a finite subcover \mathcal{U}' . This yields $B = \bigcup \mathcal{U}' \setminus \{B^c\}$, with $\mathcal{U}' \setminus \{B^c\}$ finite, thereby showing compactness of B . Thus, the Frith frame $\mathbf{\Omega}(X, \mathcal{B}) = (\Omega_{\mathcal{B}}(X), \mathcal{B})$ is coherent. Conversely, suppose that $\mathbf{\Omega}(X, \mathcal{B})$ is coherent. In particular, its underlying frame $\Omega_{\mathcal{B}}(X)$ is compact. But this is precisely the frame of opens of the patch topology of $\mathbf{Sk}_{\text{Pervin}}(X, \mathcal{B})$. Thus, the latter is compact as well. \square

As a consequence, we immediately obtain that, under the Prime Ideal Theorem, every T_0 complete Pervin space is strongly exact. We do not know whether this is a necessary hypothesis.

Corollary 5.9 *If the Prime Ideal Theorem holds, then T_0 complete Pervin spaces are strongly exact.*

Proof Let (X, \mathcal{S}) be a Pervin space. By Lemma 5.8, we only need to show that if $\mathbf{Sk}_{\text{Pervin}}(X, \mathcal{S})$ is compact then (X, \mathcal{S}) is strongly exact. Let $\bigcap_{i \in I} S_i$ be an open intersection of \mathcal{S} , say $\bigcap_{i \in I} S_i = \bigcup_{j \in J} S'_j$, for some $\{S_j\}_{j \in J} \subseteq \mathcal{S}$. Then, we have

$$X = \left(\bigcap_{i \in I} S_i \right)^c \cup \left(\bigcap_{i \in I} S_i \right) = \bigcup_{i \in I} S_i^c \cup \bigcup_{j \in J} S'_j$$

and, by compactness of $\mathbf{Sk}_{\text{Pervin}}(X, \mathcal{S})$, it follows that there exists a finite subset $J' \subseteq J$ such that $X = \bigcup_{i \in I} S_i^c \cup \bigcup_{j \in J'} S'_j$. But then, we have $\bigcap_{i \in I} S_i = \bigcup_{j \in J'} S'_j$, and thus, the intersection $\bigcap_{i \in I} S_i$ belongs to \mathcal{S} , as required. \square

It only remains to show the inclusion $\mathbf{Sk}_{\text{Pervin}}[\mathbf{CPervin}] \supseteq \mathbf{BiTop}_{\text{KZ}}$.

Lemma 5.10 *If $\mathcal{X} = (X, \tau_+, \tau_-)$ is a compact bitopological space, then the Pervin space $\mathbf{Clop}_+(\mathcal{X})$ is complete. In particular, every T_0 , compact, and zero-dimensional bitopological space is of the form $\mathbf{Sk}_{\text{Pervin}}(X, \mathcal{S})$, for some T_0 complete Pervin space (X, \mathcal{S}) .*

Proof We let τ denote the patch topology of \mathcal{X} , and we suppose that \mathcal{X} is compact, that is, that the space (X, τ) is compact. We let $(X, \mathcal{S}) = \mathbf{Clop}_+(\mathcal{X})$ and $F \subseteq \mathcal{P}(X)$ be a Cauchy filter. Since F is proper, it has the finite intersection property, and therefore, so does $F \cap \overline{\mathcal{S}}$ (which is a subset of $\text{Clop}(X, \tau)$). By compactness of (X, τ) it follows that the intersection $\bigcap (F \cap \overline{\mathcal{S}})$ is nonempty. Thus, by Lemma 4.2, F converges and (X, \mathcal{S}) is complete.

Finally, let $\mathcal{X} = (X, \tau_+, \tau_-)$ be a bitopological space. By Proposition 3.2, if \mathcal{X} is zero-dimensional, then it is isomorphic to $\mathbf{Sk}_{\text{Pervin}} \circ \mathbf{Clop}_+(\mathcal{X})$. If furthermore \mathcal{X} is compact then, by the first part of the claim, $\mathbf{Clop}_+(\mathcal{X})$ is complete. It is also easy to see that $\mathbf{Clop}_+(\mathcal{X})$ is T_0 provided \mathcal{X} is T_0 and zero-dimensional. Thus, if \mathcal{X} is T_0 , compact, and zero-dimensional, then $(X, \mathcal{S}) = \mathbf{Clop}_+(\mathcal{X})$ is a T_0 complete Pervin space satisfying $\mathcal{X} \cong \mathbf{Sk}_{\text{Pervin}}(X, \mathcal{S})$. \square

As already explained, we may thus derive the following:

Corollary 5.11 *If the Prime Ideal Theorem holds, the adjunction $\mathbf{Clop}_+ \dashv \mathbf{Sk}_{\text{Pervin}}$ restricts to an equivalence $\mathbf{BiTop}_{\text{KZ}} \cong \mathbf{CPervin}$.*

Let us now consider the pointfree setting. We have seen in Sect. 3.1 the existence of an idempotent adjunction $\mathbf{Sk}_{\text{Frith}} : \mathbf{Frith} \rightleftarrows \mathbf{BiFrm} : \mathbf{B}_+$. Although we were not able to describe the underlying categorical equivalence, we have shown that the fixpoints of \mathbf{Frith} are the strongly exact Frith frames and that every compact zero-dimensional biframe is a fixpoint of \mathbf{BiFrm} . Thus, we will follow the same strategy as for Pervin spaces and show that complete Frith frames are strongly exact (cf. Lemma 5.14) and the equality $\mathbf{Sk}_{\text{Frith}}[\mathbf{CFrith}] = \mathbf{BiFrm}_{\text{KZ}}$ holds (cf. Lemmas 5.15 and 5.16).

Before proceeding, we need to state a technical result. Recall that a filter $F \subseteq L$ on a frame L is *Scott-open* if whenever $D \subseteq F$ is a directed subset whose join belongs to F , the intersection $D \cap F$ is nonempty.

Proposition 5.12 ([14, 15]) *Scott-open filters are closed under strongly exact meets.*

Proof It is shown in [14, Lemma 3.4] that every Scott-open filter $F \subseteq L$ is of the form $\{x \in L \mid \Delta x \subseteq \bigvee_{a \in P} \Delta a\}$, for some subset $P \subseteq L$. In turn, filters of this form are shown in [15, Theorem 4.5] to be closed under strongly exact meets. □

Remark 5.13 In the proof of Proposition 5.12 we invoked [14, Lemma 3.4], whose proof uses ordinal induction (although no choice principles are required). A constructive alternative proof is provided in [23, Theorem 1.9].

We may now show that complete Frith frames are strongly exact.

Lemma 5.14 *Complete Frith frames are strongly exact.*

Proof Let (L, S) be a complete Frith frame and $P \subseteq S$ be such that $a = \bigwedge P$ is a strongly exact meet. Since, by Theorem 4.8, P consists of compact elements and finite meets of compact elements are compact too, the filter $F \subseteq L$ generated by P is Scott-open. By Proposition 5.12, a belongs to F . Thus, there exist $s_1, \dots, s_n \in P$ such that $a \geq s_1 \wedge \dots \wedge s_n$, and therefore also $a = s_1 \wedge \dots \wedge s_n$. This means that $a \in S$, as required. □

Lemma 5.15 *If (L, S) is a complete Frith frame, then the biframe $\mathbf{Sk}_{\text{Frith}}(L, S)$ is compact (and zero-dimensional).*

Proof By definition, (L, S) is complete if and only if so is $\mathbf{Sym}_{\text{Frith}}(L, S) = (C_S L, \bar{S})$. By Theorem 4.8, $(C_S L, \bar{S})$ is complete if and only if it is coherent. In particular, if (L, S) is complete, then $C_S L$ is compact. But, by definition, this means that $\mathbf{Sk}_{\text{Frith}}(L, S) = (C_S L, \nabla L, \Delta S)$ is compact. Finally, $\mathbf{Sk}_{\text{Frith}}(L, S)$ is clearly zero-dimensional, for every Frith frame (L, S) . □

Lemma 5.16 *If \mathcal{L} is a compact biframe, then the Frith frame $\mathbf{B}_+(\mathcal{L})$ is complete. In particular, every compact and zero-dimensional biframe is of the form $\mathbf{Sk}_{\text{Frith}}(L, S)$, for some complete Frith frame (L, S) .*

Proof Let $\mathcal{L} = (L, L_+, L_-)$ be a compact biframe and $(M, T) = \mathbf{B}_+(\mathcal{L})$ (that is, T is the lattice of bicomplemented elements of L_+ and M is the subframe of L it generates). By Theorem 4.8, $\mathbf{B}_+(\mathcal{L})$ is complete provided T consists of compact elements of M . Since M

is, by definition, a subframe of L_+ , it suffices to show that bicomplemented elements of L_+ are compact in L_+ . Let then $t \in L_+$ be bicomplemented and suppose that $t \leq \bigvee P$, for some subset $P \subseteq L_+$. Then, we have $1 = t \vee t^* \leq \bigvee P \vee t^*$ and, since L is compact, there exists a finite subset $P' \subseteq P$ such that $1 \leq \bigvee P' \vee t^*$. But this implies that $t \leq \bigvee P'$, as required.

Finally, if \mathcal{L} is compact and zero-dimensional then, by Proposition 3.8, $\mathcal{L} \cong \mathbf{Sk}_{\mathbf{Frith}} \circ \mathbf{B}_+(L)$ and, in particular, it is of the form $\mathbf{Sk}_{\mathbf{Frith}}(L, S)$ for some complete Frith frame (L, S) . \square

We may thus conclude the following:


Corollary 5.17 *The adjunction $\mathbf{Sk}_{\mathbf{Frith}} \dashv \mathbf{B}_+$ restricts to an equivalence $\mathbf{CFrith} \cong \mathbf{BFrm}_{\mathbf{KZ}}$.*

Corollary 5.18 *If the Prime Ideal Theorem holds, then the duality between compact zero-dimensional bitopological spaces and compact zero-dimensional biframes may be seen as restriction and co-restriction of the dual adjunction $\Omega : \mathbf{Pervin} \rightleftharpoons \mathbf{Frith} : \mathbf{pt}$ along full subcategory embeddings.*

We remark that, in [17], the duality between compact zero-dimensional bispaces and compact zero-dimensional biframes is seen as a restriction and co-restriction of the classical dual adjunction $\Omega_b : \mathbf{BiTop} \rightleftharpoons \mathbf{BFrm} : \mathbf{pt}_b$. However, although bitopological spaces and biframes are related to Pervin spaces and Frith frames (namely, via the functors $\mathbf{Sk}_{\mathbf{Pervin}}$ and $\mathbf{Sk}_{\mathbf{Frith}}$, respectively), as seen in Example 3.11, the two extensions of the adjunction $\Omega : \mathbf{Top} \rightleftharpoons \mathbf{Frm} : \mathbf{pt}$ are not comparable. The situation is different when we restrict to T_0 complete and to T_0 compact and zero-dimensional structures, as in that case, both $\Omega_b : \mathbf{BiTop} \rightleftharpoons \mathbf{BFrm} : \mathbf{pt}_b$ and $\Omega : \mathbf{CPervin} \rightleftharpoons \mathbf{CFrith} : \mathbf{pt}$ coincide, up to equivalence. Therefore, our duality ends up being equivalent to the duality of [17].

5.4 Summary

We summarize in the diagram below the categorical equivalences shown in Sects. 5.1–5.3, which exhibit several Stone-type dualities as restrictions of the dual adjunction between Pervin spaces and Frith frames. The results marked with * require the use of the Prime Ideal Theorem.



We highlight that the explicit equivalences between **BiTop**_{KZ} and **Priest**, and between **BiFrm**_{KZ} and **DLat** have first been studied in [17].

6 The T_D Axiom for Pervin Spaces and Locale-Based Frith Frames

In [7, Section 4.5] a Pervin equivalent of the T_D axiom for topological spaces is introduced. A Pervin space is called T_D if for every $x \in X$ there is $S \in \mathcal{S}$ such that $x \in S$ and $S \setminus \{x\} \in \mathcal{S}$. In there, the following slight variants of this definition were shown to be equivalent:

Lemma 6.1 *Let (X, \mathcal{S}) be a Pervin space. Then, the following are equivalent:*

- (a) (X, \mathcal{S}) is T_D ,
- (b) for every $x \in X$, there are distinct $S_1, S_2 \in \mathcal{S}$ such that $S_1 \setminus \{x\} = S_2 \setminus \{x\}$,
- (c) for every $x \in X$, there are $S_1, S_2 \in \mathcal{S}$ such that $S_1 \setminus S_2 = \{x\}$.

On the other hand, in [5] the T_D axiom is compared to sobriety and it is highlighted that the two axioms are, in some sense, duals of each other. In particular, the following is proven.

Proposition 6.2 ([5, Proposition 4.3]) *Let X be a topological space. In the category of T_0 topological spaces we have the following:*

- (a) The space X is sober if and only if whenever we have an extremal monomorphism $f : X \hookrightarrow Y$ such that $\Omega(f)$ is an isomorphism f must be an isomorphism.
- (b) The space X is T_D if and only if whenever we have an extremal monomorphism $f : Y \hookrightarrow X$ such that $\Omega(f)$ is an isomorphism f must be an isomorphism.

We will now show a Pervin version of this proposition, where *sober* is replaced by *complete* and the T_D -axiom is replaced by its equivalent for Pervin spaces, being the next lemma the key ingredient. Recall the forgetful function $\mathbf{L}_{\text{Pervin}} : \mathbf{Pervin} \rightarrow \mathbf{DLat}^{op}$ introduced in Sect. 2.6.

Lemma 6.3 *Let $f : (X, \mathcal{S}) \rightarrow (Y, \mathcal{T})$ be a morphism of Pervin spaces. Then,*

- (a) $\mathbf{L}_{\text{Pervin}}(f) : \mathcal{T} \rightarrow \mathcal{S}$ is injective if and only if $\mathbf{Sym}_{\text{Perv}}(f) : (X, \overline{\mathcal{S}}) \rightarrow (Y, \overline{\mathcal{T}})$ is dense;
- (b) $\mathbf{L}_{\text{Pervin}}(f) : \mathcal{T} \rightarrow \mathcal{S}$ is surjective if and only if $\Omega(f) : (\Omega_{\mathcal{T}}(Y), \mathcal{T}) \rightarrow (\Omega_{\mathcal{S}}(X), \mathcal{S})$ is an extremal epimorphism of Frith frames. Moreover, if (X, \mathcal{S}) is T_0 , these are further equivalent to f being an extremal monomorphism of Pervin spaces.

Proof Let us start by proving (a). Since $\mathbf{L}_{\text{Pervin}}(f)$ is a lattice homomorphism, being injective is equivalent to having

$$\forall T_1, T_2 \in \mathcal{T}, f^{-1}(T_1) \subseteq f^{-1}(T_2) \implies T_1 \subseteq T_2,$$

which is easily seen to be equivalent to having

$$\forall T_1, T_2 \in \mathcal{T}, f^{-1}(T_1 \cap T_2^c) = \emptyset \implies T_1 \cap T_2^c = \emptyset.$$

Since $\overline{\mathcal{T}}$ consists of the finite joins of elements of the form $T_1 \cap T_2^c$, with $T_1, T_2 \in \mathcal{T}$, this means that $\mathbf{Sym}_{\text{Perv}}(f)$ is dense, as required.

Finally, the first assertion of (b) is a trivial consequence of the characterization of extremal epimorphisms of Frith frames (recall Proposition 2.4), while the second one is the content of [7, Corollary 4.17]. □

Proposition 6.4 *Let (X, S) and (Y, T) be T_0 Pervin spaces. We have the following.*

- (a) *The space (X, S) is complete if and only if whenever there is a map $f : (X, S) \rightarrow (Y, T)$ such that $\mathbf{L}_{\text{Pervin}}(f)$ is an isomorphism f must be an isomorphism.*
- (b) *The space (X, S) is T_D if and only if whenever there is a map $f : (Y, T) \rightarrow (X, S)$ such that $\mathbf{L}_{\text{Pervin}}(f)$ is an isomorphism f must be an isomorphism.*

Proof Since (X, S) is T_0 , by Lemma 6.3, $\mathbf{L}_{\text{Pervin}}(f)$ is an isomorphism if and only if f is an extremal monomorphism and $\mathbf{Sym}_{\text{Perv}}(f)$ is dense. Thus, part (a) follows from Theorem 4.6(c).

To prove (b), let us first suppose that (X, S) is T_D and let $f : (Y, T) \rightarrow (X, S)$ be such that (Y, T) is T_0 and $\mathbf{L}_{\text{Pervin}}(f)$ is an isomorphism. Since (Y, T) is T_0 , by Lemma 6.3(b), f is an extremal monomorphism. Thus, it suffices to show that f is an epimorphism, that is, surjective. Fix $x \in X$. Since (X, S) is T_D , by Lemma 6.1(c), the singleton $\{x\}$ belongs to \bar{S} . Since, by Lemma 6.3(a), $\mathbf{Sym}_{\text{Perv}}(f)$ is dense, it follows that $\{x\}$ intersects $f[Y]$ (recall Lemma 2.9), that is, x belongs to the image of f as we intended to show. Conversely, let $f : (Y, T) \hookrightarrow (X, S)$ be the extremal monomorphism induced by $Y := X \setminus \{x\}$, that is, $T = \{S \setminus \{x\} \mid S \in S\}$. Clearly, (Y, T) is T_0 and f is not an isomorphism. Thus, by assumption, $\mathbf{L}_{\text{Pervin}}(f)$ cannot be injective. That is to say that, there exist distinct $S_1, S_2 \in S$ such that $S_1 \setminus \{x\} = S_2 \setminus \{x\}$ and thus, by Lemma 6.1(b), (X, S) is T_D . □

We finish this section by exhibiting a pointfree version of Proposition 6.4, which will now involve the forgetful functor $\mathbf{L}_{\text{Frith}} : \mathbf{Frith} \rightarrow \mathbf{DLat}$ (cf. Sect. 2.6). Let us start with a version of Lemma 6.3 for Frith frames.

Lemma 6.5 *Let $h : (L, S) \rightarrow (M, T)$ be a homomorphism of Frith frames. Then,*

- (a) *$\mathbf{L}_{\text{Frith}}(h) : S \rightarrow T$ is injective if and only if $\mathbf{Sym}_{\text{Frith}}(h) : (C_S L, \bar{S}) \rightarrow (C_T M, \bar{T})$ is dense;*
- (b) *$\mathbf{L}_{\text{Frith}}(h) : S \rightarrow T$ is surjective if and only if h is an extremal epimorphism.*

Proof By definition of $\mathbf{Sym}_{\text{Frith}}$, we have that $\mathbf{Sym}_{\text{Frith}}(h)$ is dense if and only if for every $s_1, s_2 \in S$, having $\nabla_{h(s_1)} \wedge \Delta_{h(s_2)} = 0$ implies that $\nabla_{s_1} \wedge \Delta_{s_2} = 0$. But this is easily seen to be equivalent to having that $h(s_1) \leq h(s_2)$ implies that $s_1 \leq s_2$, that is to say that $\mathbf{L}_{\text{Frith}}(h)$ is injective. This proves (a). Part (b) is a straightforward consequence of the definition of $\mathbf{L}_{\text{Pervin}}(h)$ and of the characterization of extremal epimorphisms of Frith frames (cf. Proposition 2.4). □

We may then prove the following version of Proposition 6.4(a). In particular, note that it provides an alternative criterion for a Frith frame to be complete, which does not depend on its symmetrization.

Proposition 6.6 *A Frith frame (L, S) is complete if and only if for every map $h : (M, T) \rightarrow (L, S)$ such that $\mathbf{L}_{\text{Frith}}(h)$ is an isomorphism, h is an isomorphism.*

Proof We first observe that (L, S) is complete if and only if every dense extremal epimorphisms $(K, C) \rightarrow (C_S L, \bar{S})$, with (K, C) symmetric, is an isomorphism. Suppose that (L, S) is complete and let $h : (M, T) \rightarrow (L, S)$ be such that $\mathbf{L}_{\text{Frith}}(h)$ is an isomorphism. By Lemma 6.5(b), h is an extremal epimorphism. Thus, we only need to show that it is also a monomorphism, that is, injective. Consider the homomorphism $\mathbf{Sym}_{\text{Frith}}(h) :$

$(C_T M, \overline{T}) \rightarrow (C_S L, \overline{S})$. Then, $\mathbf{Sym}_{\mathbf{Frith}}(h)$ is an extremal epimorphism because so is h and, by Lemma 6.5(a), it is dense. Thus, since (L, S) is complete, it is an isomorphism and, in particular, injective. Thus, its restriction h is also injective as required. Conversely, let $h : (K, C) \rightarrow (C_S L, \overline{S})$ be a dense extremal epimorphism, with (K, C) symmetric. Since h is already a morphism in $\mathbf{Frith}_{\text{sym}}$, and thus, $\mathbf{Sym}_{\mathbf{Frith}}(h) = h$, by Lemma 6.5, $\mathbf{L}_{\mathbf{Frith}}(h)$ is an isomorphism. Thus, by hypothesis, h is an isomorphism, which proves that (L, S) is complete. \square

In order to get an analogue of Proposition 6.4(b), we need to introduce the notion of *locale-based* Frith frame, which will replace the T_D axiom in our statement.

We say that a Frith frame (L, S) is *locale-based* if the smallest sublocale of L that contains S is the whole frame L . It is not hard to see that such sublocale consists of the arbitrary meets of elements of the form $a \rightarrow s$, where $a \in L$ and $s \in S$. As a consequence, we have the following characterization of locale-based Frith frames, that we state for later reference.

Lemma 6.7 *A Frith frame is locale-based if and only if, for every $a \in L$, the following equality holds:*

$$a = \bigwedge \{b \rightarrow s \mid b \in L, s \in S, \text{ and } a \leq b \rightarrow s\}.$$

The following technical result will also be useful:

Lemma 6.8 *Let $h : (L, S) \rightarrow (M, T)$ be a morphism of Frith frames. If $\mathbf{L}_{\mathbf{Frith}}(h)$ is injective, then $h_* \circ h(s) = s$, for every $s \in S$.*

Proof Since h_* is right adjoint to h , we have $h \circ h_* \circ h = h$. Thus, the claim follows from injectivity of $\mathbf{L}_{\mathbf{Frith}}(h)$. \square

We are now ready to show the already announced analogue of Proposition 6.4(b).

Proposition 6.9 *A Frith frame (L, S) is locale-based if and only if for every map $h : (L, S) \rightarrow (M, T)$ such that $\mathbf{L}_{\mathbf{Frith}}(h)$ is an isomorphism, h is an isomorphism.*

Proof Let (L, S) be a locale-based Frith frame, and $h : (L, S) \rightarrow (M, T)$ be a homomorphism such that $\mathbf{L}_{\mathbf{Frith}}(h)$ is an isomorphism. By Lemma 6.5(b), we know that h is an extremal epimorphism. It remains to show it is injective or, equivalently, that $h_* \circ h(a) \leq a$, for every $a \in L$. It remains to show it is injective or, equivalently, that $h_* \circ h(a) \leq a$, for every $a \in L$. By Lemma 6.8 we know that $S \subseteq h_*[M]$. This is also a sublocale inclusion, as $S \subseteq L$ is, L being locale-based. As $h_*[M] \subseteq L$ is a sublocale then it must be L , as L is locale-based.

For the converse, suppose that (L, S) is such that all maps $h : (L, S) \rightarrow (M, T)$ such that $\mathbf{L}_{\mathbf{Frith}}(h)$ is an isomorphism are isomorphisms. We let $\langle S \rangle_{\mathbf{Loc}}$ be the smallest sublocale of L that contains S . Since $\langle S \rangle_{\mathbf{Loc}}$ is a subposet of L , each join computed in $\langle S \rangle_{\mathbf{Loc}}$ is greater than or equal to the same join computed in L . Thus, $(\langle S \rangle_{\mathbf{Loc}}, S)$ is also a Frith frame. Finally, we let $q : L \rightarrow \langle S \rangle_{\mathbf{Loc}}$ be the frame quotient defined by the sublocale embedding $\langle S \rangle_{\mathbf{Loc}} \hookrightarrow L$. Since the restriction of q to $\langle S \rangle_{\mathbf{Loc}}$ is the identity, q induces a morphism of Frith frames $q : (L, S) \rightarrow (\langle S \rangle_{\mathbf{Loc}}, S)$. Clearly we have that $\mathbf{L}_{\mathbf{Frith}}(q)$ is an isomorphism, and so, by hypothesis, h is an isomorphism. This proves that (L, S) is locale-based, as required. \square

Author Contributions Both authors contributed to all stages of the paper.

Funding Célia Borlido was partially supported by the Centre for Mathematics of the University of Coimbra - UIDB/00324/2020, funded by the Portuguese Government through FCT/MCTES. Anna Laura Suarez received

funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation program (grant agreement No.670624).

Data Availability Not applicable

Declarations

Conflict of interests None

References

1. Ball, R.N., Picado, J., Pultr, A.: Notes on exact meets and joins. *Appl. Categ. Struct.* **22**, 699–714 (2014)
2. Banaschewski, B.: Coherent frames. In *Continuous Lattices* (Berlin, Heidelberg), B. Banaschewski and R.-E. Hoffmann, Eds., Springer Berlin Heidelberg, pp. 1–11 (1981)
3. Banaschewski, B.: Universal, zero-dimensional compactifications. In *Categorical topology and its relation to analysis, algebra and combinatorics* (Prague), World Sci. Publ. Teaneck, NJ **1989**, 257–269 (1988)
4. Banaschewski, B., Brümmer, G.C., Hardie, K.A.: Biframes and bispaces. *Quaest. Math.* **6**(1–3), 13–25 (1983)
5. Banaschewski, B., Pultr, A.: Pointfree aspects of the T_D axiom of classical topology. *Quaest. Math.* **33**(3), 369–385 (2010)
6. Bezhanishvili, G., Bezhanishvili, N., Gabelaia, D., Kurz, A.: Bitopological duality for distributive lattices and Heyting algebras. *Math. Struct. Comput. Sci.* **20**(3), 359–393 (2010)
7. Borlido, C., Suarez, A. L.: A pointfree theory of Pervin spaces. *Quaestiones Mathematicae*, 1–40 (2023)
8. Cornish, W.: On H. Priestley's dual of the category of bounded distributive lattices. *Matematički Vesnik* **12**(27), 329–332 (1975)
9. Davey, B.A., Priestley, H.A.: Introduction to lattices and order. Cambridge Mathematical Textbooks, Cambridge University Press, Cambridge (1990)
10. Frith, J. (1986) *Structured frames*. PhD thesis, University of Cape Town
11. Gehrke, M., Grigorieff, S., Pin, J.-E.: A topological approach to recognition. In *Automata, languages and programming. Part II*, vol. 6199 of Lecture Notes in Computer Science Springer, Berlin, pp. 151–162 (2010)
12. Jech, T. J.: *The axiom of choice*. Studies in Logic and the Foundations of Mathematics, Vol. 75. North-Holland Publishing Co., Amsterdam-London; American Elsevier Publishing Co., Inc., New York, (1973)
13. Johnstone, P. T.: Stone spaces, vol. 3 of Cambridge studies in advanced mathematics. Cambridge University Press, (1982)
14. Johnstone, P. T.: Vietoris locales and localic semilattices. In: *Lecture Notes in Pure and Applied Mathematics* 101 (1985). Proceedings of a conference held in the University of Bremen. (1982)
15. Moshier, M.A., Pultr, A., Suarez, A.L.: Exact and strongly exact filters. *Appl. Categ. Struct.* **28**(6), 907–920 (2020)
16. Niefeld, S., Rosenthal, K.: Spatial sublocales and essential primes. *Topol. Appl.* **26**(3), 263–269 (1987)
17. Picado, J.: Join-continuous frames, Priestley's duality and biframes. *Appl. Categ. Struct.* **2**(3), 297–313 (1994)
18. Picado, J.: *Frames and Locales: Topology without points*. Springer, Basel (2011)
19. Pin, J.E.: Dual space of a lattice as the completion of a Pervin space. In: *International Conference on Relational and Algebraic Methods in Computer Science*, pp. 24–40 (2017)
20. Reilly, I.L.: Zero dimensional bitopological spaces. *Indagationes Mathematicae (Proceedings)* **76**(2), 127–131 (1973)
21. Salbany, S.: *Bitopological spaces, compactifications and completions*. Mathematical Monographs of the University of Cape Town, No. 1. University of Cape Town, Department of Mathematics, Cape Town, (1974)
22. Schauerte, A.: Biframes. PhD thesis, McMaster University, Hamilton, Ontario, (1992)
23. Vickers, S.: Constructive points of powerlocales. *Math. Proc. Cambridge Philos. Soc.* **122**(2), 207–222 (1997)
24. Wilson, J. T.: The Assembly Tower and Some Categorical and Algebraic Aspects of Frame Theory. PhD thesis, Carnegie Mellon University, (1994)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Springer Nature or its licensor (e.g. a society or other partner) holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.