

GOURSAT COMPLETIONS

Diana RODELO and Idriss TCHOFFO NGUEFEU

Abstract. We characterize categories with weak finite limits whose regular completions give rise to Goursat categories.

Keywords. regular category, projective cover, Goursat category, 3-permutable variety.

Mathematics Subject Classification (2010). 08C05, 18A35, 18B99, 18E10.

1 Introduction

The construction of the free exact category over a category with finite limits was introduced in [3]. It was later improved to the construction of the free exact category over a category with finite weak limits (*weakly lex*) in [4]. This followed from the fact that the uniqueness of the finite limits of the original category is never used in the construction; only the existence. In [4], the authors also considered the free regular category over a weakly lex one.

An important property of the free exact (or regular) construction is that such categories always have enough (regular) projectives. In fact, an exact category \mathbb{A} may be seen as the exact completion of a weakly lex category if and only if it has enough projectives. If so, then \mathbb{A} is the exact completion of any of its *projective covers*. Such a phenomenon is captured by varieties of universal algebras: they are the exact completions of their full subcategory of free algebras.

Having this link in mind, our main interest in studying this subject is to characterize projective covers of certain algebraic categories through simpler properties involving projectives and to relate those properties to the known

varietal characterizations in terms of the existence of operations of their varietal theories (when it is the case). Such kind of studies have been done for the projective covers of categories which are: Mal'tsev [11], protomodular and semi-abelian [5], (strongly) unital and subtractive [6].

The aim of this work is to obtain characterizations of the weakly lex categories whose regular completion is a Goursat (=3-permutable) category (Propositions 4.4 and 4.6). We then relate them to the existence of the quaternary operations which characterize the varieties of universal algebras which are 3-permutable (Remark 4.7).

2 Preliminaries

In this section, we briefly recall some elementary categorical notions needed in the following.

A category with finite limits is **regular** if regular epimorphisms are stable under pullbacks, and kernel pairs have coequalizers. Equivalently, any arrow $f : A \rightarrow B$ has a unique factorisation $f = ir$ (up to isomorphism), where r is a regular epimorphism and i is a monomorphism and this factorisation is pullback stable.

A **relation** R from X to Y is a subobject $\langle r_1, r_2 \rangle : R \rightarrow X \times Y$. The opposite relation of R , denoted R^o , is the relation from Y to X given by the subobject $\langle r_2, r_1 \rangle : R \rightarrow Y \times X$. A relation R from X to X is called a relation on X . We shall identify a morphism $f : X \rightarrow Y$ with the relation $\langle 1_X, f \rangle : X \rightarrow X \times Y$ and write f^o for its opposite relation. Given two relations $R \rightarrow X \times Y$ and $S \rightarrow Y \times Z$ in a regular category, we write $SR \rightarrow X \times Z$ for their relational composite. With the above notations, any relation $\langle r_1, r_2 \rangle : R \rightarrow X \times Y$ can be seen as the relational composite $r_2 r_1^o$. The following properties are well known and easy to prove (see [1] for instance); we collect them in the following lemma:

Lemma 2.1. *Let $f : X \rightarrow Y$ be an arrow in a regular category \mathbb{C} , and let $f = ir$ be its (regular epimorphism, monomorphism) factorisation. Then:*

1. $f^o f$ is the kernel pair of f , thus $1_X \leq f^o f$; moreover, $1_X = f^o f$ if and only if f is a monomorphism;

2. ff^o is (i, i) , thus $ff^o \leq 1_Y$; moreover, $ff^o = 1_Y$ if and only if f is a regular epimorphism;
3. $ff^of = f$ and $f^of f^o = f^o$.

A relation R on X is **reflexive** if $1_X \leq R$, **symmetric** if $R^o \leq R$, and **transitive** if $RR \leq R$. As usual, a relation R on X is an **equivalence relation** when it is reflexive, symmetric and transitive. In particular, a kernel pair $\langle f_1, f_2 \rangle : \text{Eq}(f) \rightarrow X \times X$ of a morphism $f : X \rightarrow Y$ is an equivalence relation.

By dropping the assumption of uniqueness of the factorization in the definition of a limit, one obtains the definition of a weak limit. We call **weakly lex** a category with weak finite limits.

An object P in a category is (regular) **projective** if, for any arrow $f : P \rightarrow X$ and for any regular epimorphism $g : Y \twoheadrightarrow X$ there exists an arrow $h : P \rightarrow Y$ such that $gh = f$. We say that a full subcategory \mathbb{C} of \mathbb{A} is a **projective cover** of \mathbb{A} if two conditions are satisfied:

- any object of \mathbb{C} is regular projective in \mathbb{A} ;
- for any object X in \mathbb{A} , there exists a (\mathbb{C}) -cover of X , that is an object C in \mathbb{C} and a regular epimorphism $C \twoheadrightarrow X$.

When \mathbb{A} admits a projective cover, one says that \mathbb{A} has *enough projectives*.

Remark 2.2. If \mathbb{C} is a projective cover of a weakly lex category \mathbb{A} , then \mathbb{C} is also weakly lex [4]. For example, let X and Y be objects in \mathbb{C} and $X \leftarrow W \rightarrow Y$ a weak product of X and Y in \mathbb{A} . Then, for any cover $\bar{W} \twoheadrightarrow W$ of W , $X \leftarrow \bar{W} \rightarrow Y$ is a weak product of X and Y in \mathbb{C} . Furthermore, if \mathbb{A} is a regular category, then the induced morphism $W \twoheadrightarrow X \times Y$ is a regular epimorphism. Similar remarks apply to all weak finite limits.

3 Goursat categories

In this section we review the notion of Goursat category and the characterizations of Goursat categories through regular images of equivalence relations and through Goursat pushouts.

Definition 3.1. [2, 1] A regular category \mathbb{C} is called a **Goursat category** when the equivalence relations in \mathbb{C} are 3-permutable, i.e. $RSR = SRS$ for any pair of equivalence relations R and S on the same object.

When \mathbb{C} is a regular category, (R, r_1, r_2) is an equivalence relation on X and $f : X \rightarrow Y$ is a regular epimorphism, we define the **regular image of R along f** to be the relation $f(R)$ on Y induced by the (regular epimorphism, monomorphism) factorization $\langle s_1, s_2 \rangle \psi$ of the composite $(f \times f)\langle r_1, r_2 \rangle$:

$$\begin{array}{ccc} R & \xrightarrow{\psi} & f(R) \\ \langle r_1, r_2 \rangle \downarrow & & \downarrow \langle s_1, s_2 \rangle \\ X \times X & \xrightarrow{f \times f} & Y \times Y. \end{array}$$

Note that the regular image $f(R)$ can be obtained as the relational composite $f(R) = fRf^o = fr_2r_1^o f^o$. When R is an equivalence relation, $f(R)$ is also reflexive and symmetric. In a general regular category $f(R)$ is not necessarily an equivalence relation. This is the case in a *Goursat category* according to the following theorem.

Theorem 3.2. [1] A regular category \mathbb{C} is a Goursat category if and only if for any regular epimorphism $f : X \rightarrow Y$ and any equivalence relation R on X , the regular image $f(R) = fRf^o$ of R along f is an equivalence relation.

Goursat categories are well known in Universal Algebra. In fact, by a classical theorem in [10], a variety of universal algebras is a Goursat category precisely when its theory has two quaternary operations p and q such that the identities $p(x, y, y, z) = x$, $q(x, y, y, z) = z$ and $p(x, x, y, y) = q(x, x, y, y)$ hold. Accordingly, the varieties of groups, Heyting algebras and implication algebras are Goursat categories. The category of topological group, Hausdorff groups, right complemented semi-group are also Goursat categories.

There are many known characterizations of Goursat categories (see [1, 7, 8, 9] for instance). In particular the following characterization, through Goursat pushouts, will be useful:

Theorem 3.3. [7] Let \mathbb{C} be a regular category. The following conditions are equivalent:

- (i) \mathbb{C} is a Goursat category;
- (ii) any commutative diagram of type (I) in \mathbb{C} , where α and β are regular epimorphisms and f and g are split epimorphisms

$$\begin{array}{ccc}
 X & \xrightarrow{\alpha} & U \\
 \uparrow s & \Downarrow f & \uparrow t \\
 Y & \xrightarrow{\beta} & W,
 \end{array}
 \quad \text{(I)}
 \quad \begin{array}{l}
 g\alpha = \beta f \\
 \alpha s = t\beta
 \end{array}$$

(which is necessarily a pushout) is a **Goursat pushout**: the morphism $\lambda : \text{Eq}(f) \rightarrow \text{Eq}(g)$, induced by the universal property of kernel pair $\text{Eq}(g)$ of g , is a regular epimorphism.

Remark 3.4. Diagram (I) is a Goursat pushout precisely when the regular image of $\text{Eq}(f)$ along α is (isomorphic to) $\text{Eq}(g)$. From Theorem 3.3, it then follows that a regular category \mathbb{C} is a Goursat category if and only if for any commutative diagram of type (I) one has $\alpha(\text{Eq}(f)) = \text{Eq}(g)$.

Note that Theorem 3.2 characterizes Goursat categories through the property that regular images of equivalence relations are equivalence relations, while Theorem 3.3 characterizes them through the property that regular images of certain kernel pairs are kernel pairs.

4 Projective covers of Goursat categories

In this section, we characterize the categories with weak finite limits whose regular completion are Goursat categories.

Definition 4.1. Let \mathbb{C} be a weakly lex category:

1. a **pseudo-relation** on an object X of \mathbb{C} is a pair of parallel arrows $R \begin{array}{c} \xrightarrow{r_1} \\ \xrightarrow{r_2} \end{array} X$; a pseudo-relation is a relation if r_1 and r_2 are jointly monomorphic;
2. a pseudo-relation $R \begin{array}{c} \xrightarrow{r_1} \\ \xrightarrow{r_2} \end{array} X$ on X is said to be:

- **reflexive** when there is an arrow $r : X \longrightarrow R$ such that $r_1 r = 1_X = r_2 r$;
- **symmetric** when there is an arrow $\sigma : R \longrightarrow X$ such that $r_2 = r_1 \sigma$ and $r_1 = r_2 \sigma$;
- **transitive** if by considering a weak pullback

$$\begin{array}{ccc} W & \xrightarrow{p_2} & R \\ p_1 \downarrow & & \downarrow r_1 \\ R & \xrightarrow{r_2} & X, \end{array}$$

there is an arrow $t : W \longrightarrow R$ such that $r_1 t = r_1 p_1$ and $r_2 t = r_2 p_2$.

- a **pseudo-equivalence relation** if it is reflexive, symmetric and transitive.

Remark that the transitivity of a pseudo-relation $R \begin{smallmatrix} \xrightarrow{r_1} \\ \xrightarrow{r_2} \end{smallmatrix} X$ does not depend on the choice of the weak pullback of r_1 and r_2 ; in fact, if

$$\begin{array}{ccc} \bar{W} & \xrightarrow{\bar{p}_2} & R \\ \bar{p}_1 \downarrow & & \downarrow r_1 \\ R & \xrightarrow{r_2} & X, \end{array}$$

is another weak pullback, the factorization $\bar{W} \longrightarrow W$ composed with the transitivity $t : W \longrightarrow R$ ensures that the pseudo-relation is transitive also with respect to the second weak pullback.

The following property from [12] (Proposition 1.1.9) will be useful in the sequel:

Proposition 4.2. [12] *Let \mathbb{C} be a projective cover of a regular category \mathbb{A} . Let $R \begin{smallmatrix} \xrightarrow{r_1} \\ \xrightarrow{r_2} \end{smallmatrix} X$ be a pseudo-relation in \mathbb{C} and consider its (regular epimor-*

phism, monomorphism) factorization in \mathbb{A}

$$\begin{array}{ccc}
 R & \xrightarrow{(r_1, r_2)} & X \times X \\
 \searrow p & & \nearrow (e_1, e_2) \\
 & E &
 \end{array}$$

Then, R is a pseudo-equivalence relation in \mathbb{C} if and only if S is an equivalence relation in \mathbb{A} .

Definition 4.3. Let \mathbb{C} be a weakly lex category. We call \mathbb{C} a **weak Goursat category** if, for any pseudo-equivalence relation $R \xrightarrow[r_2]{r_1} X$ and any regular epimorphism $f : X \twoheadrightarrow Y$, the composite $R \xrightarrow[f r_2]{f r_1} X$ is also a pseudo-equivalence relation.

We use Remark 2.2 repeatedly in the next results.

Proposition 4.4. Let \mathbb{C} be a projective cover of a regular category \mathbb{A} . Then \mathbb{A} is a Goursat category if and only if \mathbb{C} is a weak Goursat category.

Proof. Since \mathbb{C} is a projective cover of a regular category \mathbb{A} , then \mathbb{C} is weakly lex.

Suppose that \mathbb{A} is a Goursat category. Let $R \xrightarrow[r_2]{r_1} X$ be a pseudo-equivalence relation in \mathbb{C} and let $f : X \twoheadrightarrow Y$ be a regular epimorphism in \mathbb{C} . For the (regular epimorphism, monomorphism) factorizations of $\langle r_1, r_2 \rangle$ and $\langle f r_1, f r_2 \rangle$ we get the following diagram

$$\begin{array}{ccc}
 R & \xrightarrow{\langle r_1, r_2 \rangle} & X \times X \\
 \parallel & \searrow p & \nearrow \langle e_1, e_2 \rangle \\
 & E & \\
 & \vdots w & \\
 & S & \\
 & \nearrow q & \searrow \langle s_1, s_2 \rangle \\
 R & \xrightarrow{\langle f r_1, f r_2 \rangle} & Y \times Y,
 \end{array}
 \tag{1}$$

where $w : E \rightarrow S$ is induced by the strong epimorphism p

$$\begin{array}{ccc}
 R & \xrightarrow{p} & E \\
 q \downarrow & \searrow w & \downarrow (f \times f) \langle e_1, e_2 \rangle \\
 S & \xrightarrow{\langle s_1, s_2 \rangle} & Y \times Y.
 \end{array}$$

Then w is a regular epimorphism and by the commutativity of the right side of (1), one has $S = f(E)$. By Proposition 4.2, we know that E is an equivalence relation in \mathbb{A} . Since \mathbb{A} is a Goursat category, then $S = f(E)$ is also an equivalence relation in \mathbb{A} and by Proposition 4.2, we can conclude that

$$R \begin{array}{c} \xrightarrow{fr_1} \\ \xrightarrow{fr_2} \end{array} X \text{ is a pseudo-equivalence relation in } \mathbb{C}.$$

Conversely, suppose that \mathbb{C} is a weak Goursat category. Let $R \begin{array}{c} \xrightarrow{r_1} \\ \xrightarrow{r_2} \end{array} X$ be an equivalence relation in \mathbb{A} and $f : X \rightarrow Y$ a regular epimorphism. We are going to show that $f(R) = S$

$$\begin{array}{ccc}
 R & \xrightarrow{h} & f(R) = S \\
 r_1 \downarrow & & s_1 \downarrow \\
 r_2 \downarrow & & s_2 \downarrow \\
 X & \xrightarrow{f} & Y
 \end{array}$$

is an equivalence relation; it is obviously reflexive and symmetric. In order to conclude that \mathbb{A} is a Goursat category, we must prove that S is transitive, i.e that S is an equivalence relation.

We begin by covering the regular epimorphism f in \mathbb{A} with a regular epimorphism \bar{f} in \mathbb{C} . For that we take the cover $y : \bar{Y} \rightarrow Y$, consider the pullback of y and f in \mathbb{A} and take its cover $\alpha : \bar{X} \rightarrow X \times_Y \bar{Y}$

$$\begin{array}{ccc}
 \bar{X} & \xrightarrow{\bar{f}} & \bar{Y} \\
 \alpha \searrow & & \downarrow y \\
 X \times_Y \bar{Y} & \xrightarrow{f'} & \bar{Y} \\
 x \searrow & & \downarrow y \\
 X & \xrightarrow{f} & Y
 \end{array}$$

Note that the above outer diagram is a *regular pushout*, so that

$$f \circ y = x \bar{f} \circ \quad \text{and} \quad y \circ f = \bar{f} \circ x \quad (2)$$

(Proposition 2.1 in [1]).

Next, we take the inverse image $x^{-1}(R)$ in \mathbb{A} , which in an equivalence relation since R is, and cover it to obtain a pseudo-equivalence $W \rightrightarrows \bar{X}$ in \mathbb{C} . By assumption $W \rightrightarrows \bar{X} \xrightarrow{\bar{f}} \bar{Y}$ is a pseudo-equivalence relation in \mathbb{C} so it factors through an equivalence relation, say $V \xrightarrow[v_2]{v_1} \bar{Y}$, in \mathbb{A} . We have

$$\begin{array}{ccccc}
 W & \xlongequal{\quad} & W & & \\
 w \downarrow & & \downarrow & \searrow v & \\
 x^{-1}(R) & \xrightarrow{\quad \gamma \quad} & V & & \\
 \langle \rho_1, \rho_2 \rangle \downarrow & \nearrow \pi_R & \downarrow \langle v_1, v_2 \rangle & \searrow \lambda & \\
 \bar{X} \times \bar{X} & \xrightarrow{\quad h \quad} & R & \xrightarrow{\quad} & S \\
 \langle r_1, r_2 \rangle \downarrow & \nearrow \pi_R & \downarrow \langle r_1, r_2 \rangle & \searrow \lambda & \\
 \bar{X} \times \bar{X} & \xrightarrow{\quad \bar{f} \times \bar{f} \quad} & \bar{Y} \times \bar{Y} & \xrightarrow{\quad y \times y \quad} & Y \times Y \\
 x \times x \downarrow & \nearrow \pi_R & \downarrow \langle x_1, x_2 \rangle & \searrow \lambda & \\
 X \times X & \xrightarrow{\quad f \times f \quad} & Y \times Y & & \\
 & & \downarrow \langle s_1, s_2 \rangle & &
 \end{array}$$

where γ and λ are induced by the strong epimorphisms w and v , respectively

$$\begin{array}{ccc}
 W & \xrightarrow{w} & x^{-1}(R) \\
 v \downarrow & \nearrow \gamma & \downarrow \langle \rho_1, \rho_2 \rangle \\
 V & \xrightarrow{\langle v_1, v_2 \rangle} & \bar{Y} \times \bar{Y} \\
 & & \downarrow \bar{f} \times \bar{f} \\
 & & \bar{Y} \times \bar{Y}
 \end{array}
 \quad \text{and} \quad
 \begin{array}{ccc}
 W & \xrightarrow{v} & V \\
 h\pi_R w \downarrow & \nearrow \lambda & \downarrow \langle v_1, v_2 \rangle \\
 S & \xrightarrow{\langle s_1, s_2 \rangle} & Y \times Y \\
 & & \downarrow y \times y \\
 & & Y \times Y
 \end{array}$$

Since γ is a regular epimorphism, we have $V = \bar{f}(x^{-1}(R))$. Since λ is a regular epimorphism, we have $S = y(V)$. One also has $V = y^{-1}(S)$

because

$$\begin{aligned}
 y^{-1}(S) &= y^{\circ}Sy \\
 &= y^{\circ}f(R)y \\
 &= y^{\circ}fRf^{\circ}y \\
 &= \bar{f}x^{\circ}Rx\bar{f}^{\circ} \quad (\text{by (2)}) \\
 &= \bar{f}(x^{-1}(R)) \\
 &= V.
 \end{aligned}$$

Finally, S is transitive since

$$\begin{aligned}
 SS &= yy^{\circ}Sy y^{\circ}Sy y^{\circ} && (\text{Lemma 2.1(2)}) \\
 &= yy^{-1}(S)y^{-1}(S)y^{\circ} \\
 &= yVVy^{\circ} \\
 &= yVy^{\circ} && (\text{since } V \text{ is an equivalence relation}) \\
 &= y(V) \\
 &= S.
 \end{aligned}$$

□

We may also consider weak Goursat categories through a property which is more similar to the one mentioned in Theorem 3.2:

Lemma 4.5. *Let \mathbb{C} be a projective cover of a regular category \mathbb{A} . Then \mathbb{C} is a weak Goursat category if and only if for any commutative diagram in \mathbb{C}*

$$\begin{array}{ccc}
 R & \xrightarrow{\varphi} & S \\
 r_1 \downarrow & & \downarrow s_1 \\
 X & \xrightarrow{f} & Y \\
 & & \downarrow s_2
 \end{array} \quad (3)$$

such that f and φ are regular epimorphism and R is a pseudo-equivalence relation, then S is a pseudo-equivalence relation.

Proof. (i) \Rightarrow (ii) Since $R \xrightarrow[r_2]{r_1} X$ is a pseudo-equivalence relation, by assumption $R \xrightarrow[f r_2]{f r_1} X$ is also a pseudo-equivalence relation and then its (regular epimorphism, monomorphism) factorization gives an equivalence

relation $E \begin{smallmatrix} \xrightarrow{e_1} \\ \xrightarrow{e_2} \end{smallmatrix} Y$ in \mathbb{A} (Proposition 4.2). We have the following commutative diagram

$$\begin{array}{ccccc}
 R & \xlongequal{\quad} & R & \xrightarrow{\varphi} & S \\
 \downarrow r_1 & & \downarrow fr_1 & \searrow \rho & \downarrow \sigma \\
 & & & & E \\
 \downarrow r_2 & & \downarrow fr_2 & \nearrow e_1 & \downarrow \sigma \\
 X & \xrightarrow{f} & Y & & S \\
 & & & \nearrow e_2 & \downarrow s_1 \\
 & & & & Y \\
 & & & & \downarrow s_2
 \end{array}$$

where $\sigma : S \rightarrow E$ is induced by the strong epimorphism φ

$$\begin{array}{ccc}
 R & \xrightarrow{\varphi} & S \\
 \rho \downarrow & \searrow \sigma & \downarrow \langle s_1, s_2 \rangle \\
 E & \xrightarrow{\langle e_1, e_2 \rangle} & Y \times Y.
 \end{array}$$

Then σ is a regular epimorphism and $S \begin{smallmatrix} \xrightarrow{s_1} \\ \xrightarrow{s_2} \end{smallmatrix} Y$ is a pseudo-equivalence relation (Proposition 4.2).

(ii) \Rightarrow (i) Let $R \begin{smallmatrix} \xrightarrow{r_1} \\ \xrightarrow{r_2} \end{smallmatrix} X$ be a pseudo-equivalence relation in \mathbb{C} and $f : X \rightarrow Y$ a regular epimorphism. The following diagram is of the type (3)

$$\begin{array}{ccc}
 R & \xlongequal{\quad} & R \\
 \downarrow r_1 & & \downarrow fr_1 \\
 & & \downarrow r_2 & & \downarrow fr_2 \\
 X & \xrightarrow{f} & Y.
 \end{array}$$

Since $R \begin{smallmatrix} \xrightarrow{r_1} \\ \xrightarrow{r_2} \end{smallmatrix} X$ is a pseudo-equivalence relation, then by assumption

$R \begin{smallmatrix} \xrightarrow{fr_1} \\ \xrightarrow{fr_2} \end{smallmatrix} Y$ is also a pseudo-equivalence relation. □

Alternatively, weak Goursat categories may be characterized through a property more similar to the one mentioned in Remark 3.4:

Proposition 4.6. *Let \mathbb{C} be a projective cover of a regular category \mathbb{A} . The following conditions are equivalent:*

- (i) \mathbb{A} is a Goursat category;
- (ii) \mathbb{C} is a weak Goursat category;
- (iii) For any commutative diagram of type (I) in \mathbb{C} where

$$\begin{array}{ccc}
 F & \xrightarrow{\lambda} & G \\
 \beta_1 \downarrow & & \downarrow \rho_1 \\
 X & \xrightarrow{\alpha} & U \\
 s \uparrow & & \uparrow t \\
 Y & \xrightarrow{\beta} & W
 \end{array}
 \quad (I)$$

F is a weak kernel pair of f and λ is a regular epimorphism (in \mathbb{C}), then G is a weak kernel pair of g .

Proof. (i) \Leftrightarrow (ii) By Proposition 4.4.
 (i) \Rightarrow (iii) If we take the kernel pairs of f and g , then the induced morphism $\bar{\alpha} : \text{Eq}(f) \rightarrow \text{Eq}(g)$ is a regular epimorphism by Theorem 3.3. Moreover, the induced morphism $\varphi : F \rightarrow \text{Eq}(f)$ is also a regular epimorphism. We get

$$\begin{array}{ccc}
 F & \xrightarrow{\lambda} & G \\
 \downarrow \varphi & & \downarrow w \\
 \text{Eq}(f) & \xrightarrow{\bar{\alpha}} & \text{Eq}(g) \\
 f_1 \downarrow & & \downarrow g_1 \\
 X & \xrightarrow{\alpha} & U \\
 s \uparrow & & \uparrow t \\
 Y & \xrightarrow{\beta} & W,
 \end{array}$$

$\rho_1 \swarrow \searrow \rho_2$

where $w : G \rightarrow \text{Eq}(g)$ is induced by the strong epimorphism λ

$$\begin{array}{ccc}
 F & \xrightarrow{\lambda} & G \\
 \bar{\alpha} \cdot \varphi \downarrow & & \downarrow \langle \rho_1, \rho_2 \rangle \\
 \text{Eq}(g) & \xrightarrow{\langle g_1, g_2 \rangle} & U.
 \end{array}$$

This implies that ω is a regular epimorphism ($\omega\lambda = \bar{\alpha}\varphi$) and then $G \begin{smallmatrix} \xrightarrow{\rho_1} \\ \xrightarrow{\rho_2} \end{smallmatrix} U$ is a weak kernel pair of g .

(iii) \Rightarrow (ii) Consider the diagram (3) in \mathbb{C} where $R \begin{smallmatrix} \xrightarrow{r_1} \\ \xrightarrow{r_2} \end{smallmatrix} X$ is a pseudo-equivalence relation. We want to prove that $S \begin{smallmatrix} \xrightarrow{s_1} \\ \xrightarrow{s_2} \end{smallmatrix} Y$ is also a pseudo-equivalence. Take the (regular epimorphism, monomorphism) factorization of R and S in \mathbb{A} and the induced morphism μ making the following diagram commutative

$$\begin{array}{ccc}
 R & \xrightarrow{\varphi} & S \\
 \downarrow \rho & & \downarrow \sigma \\
 U & \xrightarrow{\mu} & V \\
 \downarrow u_1 & & \downarrow v_1 \\
 X & \xrightarrow{f} & Y \\
 \downarrow u_2 & & \downarrow v_2
 \end{array}$$

Since μ is a regular epimorphism, then $V = f(U)$ and consequently, V is reflexive and symmetric.

Since S is a pseudo-relation associated to V , then S is also a reflexive and symmetric pseudo-relation. We just need to prove that V is transitive. To do so, we apply our assumption to the diagram

$$\begin{array}{ccc}
 F & \xrightarrow{\lambda} & G \\
 \downarrow \delta & \searrow & \downarrow \alpha \\
 \text{Eq}(r_1) & \xrightarrow{\chi} & \varphi(\text{Eq}(r_1)) \\
 \downarrow & & \downarrow \\
 R & \xrightarrow{\varphi} & S \\
 \downarrow e_R & & \downarrow e_S \\
 X & \xrightarrow{f} & Y
 \end{array}$$

where G is a cover of the regular image $\varphi(\text{Eq}(r_1))$ and F is a cover of the pullback $\text{Eq}(r_1) \times_{\varphi(\text{Eq}(r_1))} G$. Since δ is a regular epimorphism, then

$F \rightrightarrows R$ is a weak kernel pair of r_1 . By assumption $G \rightrightarrows S$ is a weak kernel pair of s_1 , thus $\varphi(\text{Eq}(r_1)) = \text{Eq}(s_1)$. We then have

$$\begin{aligned}
 VV &= v_2 v_1^o v_1 v_2^o && \text{(since } V \text{ is symmetric)} \\
 &= v_2 \sigma \sigma^o v_1^o v_1 \sigma \sigma^o v_2^o && \text{(Lemma 2.1(2))} \\
 &= s_2 s_1^o s_1 s_2^o && (v_i \sigma = s_i) \\
 &= s_2 \varphi r_1^o r_1 \varphi^o s_2^o && (\varphi(\text{Eq}(r_1)) = \text{Eq}(s_1)) \\
 &= f r_2 r_1^o r_1 r_2^o f^o && (s_i \varphi = f r_i) \\
 &= f u_2 \rho \rho^o u_1^o u_1 \rho \rho^o u_2^o f^o && (u_i \rho = r_i) \\
 &= f u_2 u_1^o u_1 u_2^o f^o && \text{(Lemma 2.1(2))} \\
 &= f U U f^o && \text{(since } U \text{ is an equivalence relation in } \mathbb{A}) \\
 &= f U f^o \\
 &= V.
 \end{aligned}$$

□

Remark 4.7. When \mathbb{A} is a 3-permutable variety and \mathbb{C} its subcategory of free algebras, then the property stated in Proposition 4.6 (iii) is precisely what is needed to obtain the existence of the quaternary operations p and q which characterize 3-permutable varieties. Let X denote the free algebra on one element. Diagram (I) below belongs to \mathbb{C}

$$\begin{array}{ccc}
 F & \xlongequal{\quad} & F \\
 \downarrow \mu & & \downarrow \lambda \mu \\
 \text{Eq}(\nabla_2 + \nabla_2) & \xrightarrow{\quad \lambda \quad} & \text{Eq}(\nabla_3) \\
 \begin{array}{c} \pi_1 \downarrow \downarrow \pi_2 \\ \downarrow \downarrow \\ 4X \end{array} & \xrightarrow{\quad 1 + \nabla_2 + 1 \quad} & \begin{array}{c} \downarrow \downarrow \\ 3X \end{array} \\
 \begin{array}{c} \iota_2 + \iota_1 \uparrow \downarrow \\ \uparrow \downarrow \\ 2X \end{array} & \xrightarrow{\quad} & \begin{array}{c} \iota_2 \uparrow \downarrow \\ \uparrow \downarrow \\ X \end{array} \\
 & & \text{(I)}
 \end{array}$$

If F is a cover of $\text{Eq}(\nabla_2 + \nabla_2)$, then $F \rightrightarrows 4X$ is a weak kernel pair of $\nabla_2 + \nabla_2$. By assumption $F \rightrightarrows 3X$ is a weak kernel pair of ∇_3 , so that $\lambda \mu$ is surjective. We then conclude that λ is surjective and the existence of the quaternary operations p and q follows from Theorem 3 in [7].

Acknowledgements

The first author acknowledges partial financial assistance by Centro de Matemática da Universidade de Coimbra—UID/MAT/00324/2013, funded by the Portuguese Government through FCT/MCTES and co-funded by the European Regional Development Fund through the Partnership Agreement PT2020.

The second author acknowledges financial assistance by Fonds de la Recherche Scientifique-FNRS Crédit Bref Séjour à l'étranger 2018/V 3/5/033 - IB/JN - 11440, which supported his stay at the University of Algarve, where this paper was partially written.

References

- [1] A. Carboni, G.M. Kelly, M.C. Pedicchio, *Some remarks on Mal'tsev and Goursat categories*, Appl. Categ. Structures 1 (1993), no. 4, 385-421.
- [2] A. Carboni, J. Lambek, M.C. Pedicchio, *Diagram chasing in Mal'cev categories*, J. Pure Appl. Algebra 69 (1991) 271–284.
- [3] A. Carboni and R. Celia Magno, *The free exact category on a left exact one*, J. Austr. Math. Soc. Ser., A 33 (1982) 295-301.
- [4] A. Carboni and E. Vitale, *Regular and exact completions*, J. Pure Appl. Algebra **125** (1998) 79-116.
- [5] M. Gran, *Semi-abelian exact completions*, Homology, Hom. Appl. **4** (2002) 175-189.
- [6] M. Gran and D. Rodelo, *On the characterization of Jónsson-Tarski and of subtractive varieties*, Diagrammes, Suppl. Vol. 67-68 (2012) 101-116.
- [7] M. Gran and D. Rodelo, *A new characterisation of Goursat categories*, Appl. Categ. Structures 20 (2012), no. 3, 229-238.
- [8] M. Gran and D. Rodelo, *Beck-Chevalley condition and Goursat categories*, J. Pure Appl. Algebra, 221 (2017) 2445-2457.

- [9] M. Gran, D. Rodelo and I. Tchoffo Nguéfeu, *Some remarks on connecteurs and groupoids in Goursat categories*, Logical Methods in Computer Science (2017) Vol. 13(3:14), 1-12.
- [10] J. Hagemann and A. Mitschke, *On n -permutable congruences*, Algebra Universalis 3 (1973), 8-12.
- [11] J. Rosický and E.M. Vitale, *Exact completions and representations in abelian categories*, Homology, Hom. Appl. **3** (2001) 453-466.
- [12] E. Vitale, *Left covering functors*, PhD thesis, Université catholique de Louvain (1994).

D. RODELO AND I. TCHOFFO NGUEFEU GOURSAT COMPLETIONS

Diana Rodelo
CMUC, Department of Mathematics
University of Coimbra
3001–501 Coimbra, Portugal
Departamento de Matemática, Faculdade de Ciências e Tecnologia
Universidade do Algarve, Campus de Gambelas
8005–139 Faro, Portugal
drodelo@ualg.pt

Idriss Tchoffo Nguéfeu
Institut de Recherche en Mathématique et Physique
Université Catholique de Louvain
Chemin du Cyclotron 2, 1348 Louvain-la-Neuve, Belgium
idriss.tchoffo@uclouvain.be